

BIRATIONAL MAPS PRESERVING THE CONTACT STRUCTURE ON $\mathbb{P}_{\mathbb{C}}^3$

DOMINIQUE CERVEAU AND JULIE DÉSERTI

ABSTRACT. We study the group of polynomial automorphisms of $\mathbb{C}^3$ (resp. birational self-maps of $\mathbb{P}_{\mathbb{C}}^3$) that preserve the contact structure.

2010 *Mathematics Subject Classification.* — 14E05, 14E07.

1. INTRODUCTION

In this article we work on contact structures on $\mathbb{P}_{\mathbb{C}}^3$ which is justified by the following result due to Frantzen and Peternell: if M is a compact Kähler contact threefold, then either $M \simeq \mathbb{P}_{\mathbb{C}}^3$, or $M = \mathbb{P}(T_{\mathcal{S}})$ for a Kähler surface $\mathcal{S}$ (see [19]). On $\mathbb{P}_{\mathbb{C}}^3$ there is, up to automorphisms, only one (non-singular) contact structure given in homogeneous coordinates by the 1-form $\tilde{\vartheta} = z_0 dz_1 - z_1 dz_0 + z_2 dz_3 - z_3 dz_2$. In $\mathbb{C}^3$ there is the Darboux 1-form $\omega = z_0 dz_1 + dz_2$ that is the standard local model of contact forms; it thus defines a holomorphic contact structure on $\mathbb{C}^3$ that extends to $\mathbb{P}_{\mathbb{C}}^3$ meromorphically. Note that ω has poles of order 3 along the hyperplane $z_3 = 0$. We denote by $c(\omega)$ the (meromorphic) contact structure induced on $\mathbb{P}_{\mathbb{C}}^3$ by ω . Let us remark that actually ω is birationally conjugate to $\tilde{\vartheta}|_{z_3=1}$ (more precisely they are conjugate via a polynomial automorphism in the affine chart $z_3 = 1$).

The contact geometry has a long story. The Darboux local model $z_0 dz_1 + dz_2$ is related to the formalization of $z_0 = -\frac{dz_2}{dz_1}$. For instance if S is a surface in $\mathbb{C}^3$ given by $F(z_0, z_1, z_2) = 0$ then the restriction of ω to S corresponds to the implicit differential equation $F\left(-\frac{\partial z_2}{\partial z_1}, z_1, z_2\right) = 0$. A birational self-map of $\mathbb{P}_{\mathbb{C}}^3$ which preserves the contact structure (i.e., which sends the 1-form $z_0 dz_1 + dz_2$ viewed in the affine chart $z_3 = 1$ onto a multiple of $z_0 dz_1 + dz_2$ by a rational function) is said to be a contact map. The space $\mathbb{C}^3$ with the contact form ω can be seen as an affine chart of the projectivization of the cotangent bundle $T^*\mathbb{C}^2$ (equipped with the standard Liouville contact form). As a consequence there is a natural extension of any birational self-map of the (z_1, z_2) plane ([25])

$$\mathcal{K}: \text{Bir}(\mathbb{P}_{\mathbb{C}}^2) \hookrightarrow \text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}, \quad (\phi_1, \phi_2) \mapsto \left(-\frac{\partial \phi_2}{\partial z_1} + \frac{\partial \phi_2}{\partial z_2} z_0, \phi_1(z_1, z_2), \phi_2(z_1, z_2) \right)$$

where $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$ denotes the group of contact birational self-maps of $\mathbb{P}_{\mathbb{C}}^3$. The image of $\mathcal{K}$ is the Klein group $\mathcal{K}$. In 1926 Klein conjectured that the group of contact maps is generated by $\mathcal{K}$ and the Legendre involution

$$(z_0, z_1, z_2) \mapsto (z_1, z_0, -z_2 - z_0 z_1).$$

In 2008 Gizatullin proved this "conjecture" in the case in which the contact transformations are polynomial automorphisms of the affine space. A description of the group of contact polynomial automorphisms in higher dimension is also provided ([23]).

Let G be a subgroup of the group $\text{Bir}(\mathbb{P}_{\mathbb{C}}^n)$ of birational self-maps of $\mathbb{P}_{\mathbb{C}}^n$, and let β be a meromorphic p -form on $\mathbb{P}_{\mathbb{C}}^n$; denote by

$$G_{\beta} = \{\phi \in G \mid \phi^* \beta = \beta\}$$

the subgroup of elements of G that preserve the form β . In the same spirit for 1-forms β we set

$$G_{c(\beta)} = \{\phi \in G \mid \phi^* \beta \wedge \beta = 0\}.$$

We have the obvious inclusions $G_{\beta} \subset G_{c(\beta)} \subset G$.

We first describe the group $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ of polynomial automorphisms of $\mathbb{C}^3$ that preserve the contact structure:

Theorem A. *If η is the form $d\omega = dz_0 \wedge dz_1$, then*

$$\text{Aut}(\mathbb{C}^3)_{\omega} \simeq \text{Aut}(\mathbb{C}^2)_{\eta} \ltimes \mathbb{C}, \quad \text{Aut}(\mathbb{C}^3)_{c(\omega)} \simeq \text{Aut}(\mathbb{C}^3)_{\omega} \ltimes \mathbb{C}^*.$$

Hence, as Banyaga did in the context of contact diffeomorphisms of smooth real manifolds ([2, 3, 4]), one gets that the commutator of $\text{Aut}(\mathbb{C}^3)_{\omega}$ (resp. $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$) is perfect. Any automorphism of $\text{Aut}(\mathbb{C}^2)$ is the composition of an inner automorphism and an automorphism of the field $\mathbb{C}$ (see [16]). Following this idea we describe the group $\text{Aut}(\text{Aut}(\mathbb{C}^3)_{\omega})$.

Danilov and Gizatullin proved that any finite subgroup of $\text{Aut}(\mathbb{C}^2)$ is linearizable ([24]). We obtain a similar statement:

Theorem B. *Any finite subgroup of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ is linearizable via an element of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$.*

We also deal with $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$. If ϕ belongs to $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$, then $\phi^* \omega = V(\phi) \omega$ where $V(\phi)$ is some rational function. In particular one gets a map V from $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$ to the set of rational functions in z_0, z_1, z_2 satisfying cocycle conditions: $V(\phi \circ \psi) = (V(\phi) \circ \psi) \cdot V(\psi)$.

The equality $\phi^* \omega = V(\phi) \omega$ can be rewritten as the following system of P.D.E.

$$(S) \begin{cases} \phi_0 \frac{\partial \phi_1}{\partial z_0} + \frac{\partial \phi_2}{\partial z_0} = 0 & (\star_1) \\ \phi_0 \frac{\partial \phi_1}{\partial z_1} + \frac{\partial \phi_2}{\partial z_1} = V(\phi) z_0 & (\star_2) \\ \phi_0 \frac{\partial \phi_1}{\partial z_2} + \frac{\partial \phi_2}{\partial z_2} = V(\phi) & (\star_3) \end{cases}$$

The first equation $(\star_1)$ has a special family of solutions: maps for which both ϕ_1 and ϕ_2 do not depend on z_0 ; we can then compute ϕ_0 from the two other equations. Taking (ϕ_1, ϕ_2) in $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ we get by this way the group $\mathcal{K}$.

Assume now that ϕ_1 or ϕ_2 depends on z_0 then both depend on it and (S) implies the following equality

$$\frac{\frac{\partial \phi_2}{\partial z_1} - z_0 \frac{\partial \phi_2}{\partial z_2}}{\frac{\partial \phi_2}{\partial z_0}} = \frac{\frac{\partial \phi_1}{\partial z_1} - z_0 \frac{\partial \phi_1}{\partial z_2}}{\frac{\partial \phi_1}{\partial z_0}}.$$

Let us defined α the map from $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$ into the set of rational functions in z_0, z_1 and z_2 by: $\alpha(\phi) = \infty$ if ϕ belongs to $\mathcal{K}$ and

$$\alpha(\phi) = \frac{\frac{\partial \phi_2}{\partial z_1} - z_0 \frac{\partial \phi_2}{\partial z_2}}{\frac{\partial \phi_2}{\partial z_0}} = \frac{\frac{\partial \phi_1}{\partial z_1} - z_0 \frac{\partial \phi_1}{\partial z_2}}{\frac{\partial \phi_1}{\partial z_0}}$$

otherwise.

If ϕ_1 and ϕ_2 are some first integrals of the rational vector field

$$Z_{\phi} = \alpha(\phi) \frac{\partial}{\partial z_0} - \frac{\partial}{\partial z_1} + z_0 \frac{\partial}{\partial z_2},$$

one gets ϕ_0 thanks to the first equation of (S) . Such ϕ is not necessary birational but only rational; nevertheless one gets a lot of contact birational self-maps by this way. Remark that since $\mathcal{K}$ (resp. $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$) is a subgroup of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$ there is a natural left translation action of $\mathcal{K}$ (resp. $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$) on $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$. These two actions admit a complete invariant:

Theorem C. *The map α is a complete invariant of the left translation action of $\mathcal{K}$ on $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$, that is for any ϕ and ψ in $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$ one has $\alpha(\phi) = \alpha(\psi)$ if and only if $\psi\phi^{-1}$ belongs to $\mathcal{K}$.*

The map V is a complete invariant of the left translation action of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$ of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$, i.e. for any ϕ, ψ in $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$ one has $V(\phi) = V(\psi)$ if and only if $\psi\phi^{-1}$ belongs to $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$.

We prove that α is not surjective: generic linear differential equations of second order give linear functions that are not in the image of α . Painlevé equations give examples of polynomials of higher degree that do not belong to $\text{im } \alpha$. The map V is also not surjective.

Since ω has no integral surface in $\mathbb{C}^3$ a contact birational self-map ϕ either preserves the hyperplane $z_3 = 0$, or blows down $z_3 = 0$. This naturally implies the following definition: $\phi \in \text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{c(\omega)}$ is regular at infinity if $z_3 = 0$ is preserved by ϕ and if $\phi|_{z_3=0}$ is birational. One shows that

Proposition D. *The set of maps of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$ that are regular coincides with $\text{Aut}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$.*

Let $\varsigma: \text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega} \rightarrow \text{Bir}(\mathbb{P}_{\mathbb{C}}^2)_{\eta}$ be the projection onto the two first components. We say that $\phi \in \text{Bir}(\mathbb{P}_{\mathbb{C}}^2)_{\eta}$ is exact if ϕ can be lifted via ς to $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$. One establishes the following criterion:

Theorem E. *A map $\phi = (\phi_0, \phi_1) \in \text{Bir}(\mathbb{P}_{\mathbb{C}}^2)_{\eta}$ is exact if and only if the closed form $\phi_0 d\phi_1 - z_0 dz_1$ has trivial residues. In that case $\phi_0 d\phi_1 - z_0 dz_1 = -db$ with $b \in \mathbb{C}(z_0, z_1)$ and $\phi = (\phi, z_2 + b(z_0, z_1)) \in \text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$.*

We give a lot of examples, and even subgroups, of exact maps but also prove that the map ς is not surjective:

Theorem F. *A generic quadratic element of $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)_{\eta}$ is not exact.*

Furthermore we look at invariant curves and surfaces. Thanks to a local argument of contact geometry one gets that if ϕ belongs to $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$, if m is a periodic point of ϕ , and if there exists a germ of irreducible curve C invariant by ϕ and passing through m , then either C is a curve of periodic points, or C is a legendrian curve. We also give a precise description of elements of $\text{Aut}(\mathbb{C}^3)_{\omega}$ (resp. $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$) that preserve a surface.

Besides we deal with some group properties. Danilov proved that $\text{Aut}(\mathbb{C}^2)_{\eta}$ is not simple ([15]); Cantat and Lamy showed that $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ is not simple ([12]). In the same spirit we establish that

Theorem G. *The groups $\text{Aut}(\mathbb{C}^3)_{\omega}$, $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$, $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$, the derived group of $\text{Aut}(\mathbb{C}^3)_{\omega}$ and the derived group of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ are not simple.*

Lamy proved that $\text{Aut}(\mathbb{C}^2)$ satisfies the Tits alternative ([29]), then Cantat showed that $\text{Bir}(\mathbb{P}_{\mathbb{C}}^2)$ also ([?]). In our context one gets that

Theorem H. *The groups $\text{Aut}(\mathbb{C}^3)_{\omega}$, $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ and $\text{Bir}(\mathbb{P}_{\mathbb{C}}^3)_{\omega}$ satisfy the Tits alternative.*

Acknowledgments. We would like to thank Guy Casale for discussions about the non-integrability.

CONTENTS

1. Introduction	1
Part 1. Contact polynomial automorphisms	4
2. Contact forms and contact structures	4
3. Description of contact automorphisms	5
4. Automorphisms group	8
Part 2. Contact birational maps	10
5. A P.D.E. approach	11
6. Invariant forms and vector fields	15
7. Regular birational maps	17
8. Exact birational maps	18
9. Indeterminacy and exceptional sets	23
Part 3. Some common properties	25
10. Invariant curves and surfaces	25
11. Dynamical properties	28
12. Non-simplicity, Tits alternative	32
13. Non-conjugate isomorphic groups	33
Part 4. Appendix: Automorphisms group of $\text{Aut}(\mathbb{C}^2)_\eta$	33
References	34

Part 1. Contact polynomial automorphisms

A **polynomial automorphism** ϕ of $\mathbb{C}^n$ is a polynomial map of the type

$$\phi: \mathbb{C}^n \rightarrow \mathbb{C}^n, \quad (z_0, z_1, \dots, z_{n-1}) \mapsto (\phi_0(z_0, z_1, \dots, z_{n-1}), \phi_1(z_0, z_1, \dots, z_{n-1}), \dots, \phi_{n-1}(z_0, z_1, \dots, z_{n-1}))$$

that is bijective. The set of polynomial automorphisms of $\mathbb{C}^n$ form a group denoted $\text{Aut}(\mathbb{C}^n)$.

The automorphisms of $\mathbb{C}^n$ of the form $(\phi_0, \phi_1, \dots, \phi_{n-1})$ where ϕ_i depends only on $z_i, z_{i+1}, \dots, z_{n-1}$ form the **Jonquières subgroup** $J_n \subset \text{Aut}(\mathbb{C}^n)$. Moreover one has the inclusions

$$\text{GL}(\mathbb{C}^n) \subset \text{Aff}_n \subset \text{Aut}(\mathbb{C}^n)$$

where Aff_n denotes the **group of affine maps**

$$\phi: (z_0, z_1, \dots, z_{n-1}) \mapsto (\phi_0(z_0, z_1, \dots, z_{n-1}), \phi_1(z_0, z_1, \dots, z_{n-1}), \dots, \phi_{n-1}(z_0, z_1, \dots, z_{n-1}))$$

with ϕ_i affine; Aff_n is the semi-direct product of $\text{GL}(\mathbb{C}^n)$ with the commutative subgroups of translations. The subgroup $\text{Tame}_n \subset \text{Aut}(\mathbb{C}^n)$ generated by J_n and Aff_n is called the **group of tame automorphisms**.

Convention: In all the article we denote $\mathbb{P}_{\mathbb{C}}^n$ by $\mathbb{P}^n$, and we write "birational maps of $\mathbb{P}^n$ " instead of "birational self-maps of $\mathbb{P}^n$ ".

2. CONTACT FORMS AND CONTACT STRUCTURES

We recall in the context of 3-manifolds the formalism of contact structure. Let M be a complex 3-manifold; we denote by $\Omega^i(M)$ the space of holomorphic i -forms on M . A **contact form** on M is an element

$\Theta \in \Omega^1(M)$ such that the 3-form $\Theta \wedge d\Theta \in \Omega^3(M)$ has no zero: $\Theta \wedge d\Theta(m) \neq 0$ for any $m \in M$. For such a contact form there is a local model given by Darboux theorem: at each point m there is a local biholomorphism $F: M_{,m} \rightarrow \mathbb{C}_{,0}^3$ such that $\Theta = F^*(z_0 dz_1 + dz_2)$. The 1-form $z_0 dz_1 + dz_2$ is called the **standard contact form** on $\mathbb{C}^3$; we denote it by ω .

A **contact structure** on the 3-manifold M is given by the following data:

- i. an open covering $M = \sqcup_k \mathcal{U}_k$,
- ii. on each $\mathcal{U}_k$ a contact form $\Theta_k \in \Omega^1(\mathcal{U}_k)$,
- iii. on each non-trivial intersection $\mathcal{U}_k \cap \mathcal{U}_\ell$ a holomorphic unit $g_{k\ell} \in \mathcal{O}^*(\mathcal{U}_k \cap \mathcal{U}_\ell)$ such that $\Theta_k = g_{k\ell} \Theta_\ell$.

A contact structure defines a holomorphic hyperplanes field $t: M \rightarrow \mathbb{P}(\text{TM})^\vee$ given for all $m \in \mathcal{U}_k$ by

$$t(m) = \ker \Theta_k(m).$$

As we recalled in §1 the compact Kähler manifolds having a contact structure are classified by Frantzen and Peternell theorem ([19]). On $\mathbb{P}^3$ there is no contact form because there is no non-trivial global form. Nevertheless there are contact structures; one of them is given in homogeneous coordinates by the 1-form

$$\tilde{\vartheta} = z_0 dz_1 - z_1 dz_0 + z_2 dz_3 - z_3 dz_2.$$

In that case we can take the standard covering by affine charts $\mathcal{U}_k = \{z_k = 1\}$ and $\vartheta_k = \tilde{\vartheta}|_{\mathcal{U}_k}$.

Proposition 2.1. *Up to automorphisms of $\mathbb{P}^3$ there is only one contact structure on $\mathbb{P}^3$.*

Proof. Remark that to a contact structure on $\mathbb{P}^3$ is associated a homogeneous 1-form β on $\mathbb{C}^4$ such that $\mathcal{U}_k = \{z_k = 1\}$ and $\Theta_k = \beta|_{\mathcal{U}_k}$ satisfies properties i., ii., iii.

Let β be a contact structure on $\mathbb{P}^3$, and let $R = \sum_i z_i \frac{\partial}{\partial z_i}$ be the radial vector field. Since $i_R \beta = 0$, to give β is equivalent to give $d\beta$. According to [26, Chapter 2, Proposition 2.1] one has $\deg d\beta = 0$; to give $d\beta$ is thus equivalent to give an antisymmetric matrix of maximal rank. But up to conjugacy there is only one 4×4 antisymmetric matrix of maximal rank. $\square$

Remark 2.2. The group of linear automorphisms of $\mathbb{C}^4$ that preserve $\tilde{\vartheta}$ coincides with the group of automorphisms of $\mathbb{P}^3$ that preserve $d\tilde{\vartheta}$; as a consequence the subgroup of $\text{Aut}(\mathbb{P}^3)$ that preserves the contact structure associated to $d\tilde{\vartheta}$ is the projectivization of the symplectic group $\text{Sp}(4; \mathbb{C})$.

Remark that the data of a global meromorphic 1-form Θ on M such that $\Theta \wedge d\Theta \neq 0$ induces a contact form (and a contact structure) on the complement of the poles and zeros of Θ and $\Theta \wedge d\Theta$. In that case we say that Θ induces a **meromorphic contact structure** on M .

For instance the Darboux form $\omega = z_0 dz_1 + dz_2$ induces a meromorphic contact structure on $\mathbb{P}^3$. In fact the forms ω and $\tilde{\vartheta}|_{z_3=1}$ are conjugate on $\mathbb{C}^3$ via $(\frac{z_0}{2}, z_1, -z_2 + \frac{z_0 z_1}{2})$. The corresponding (meromorphic) contact structure are birationally conjugate on $\mathbb{P}^3$.

3. DESCRIPTION OF CONTACT AUTOMORPHISMS

3.1. Description of $\text{Aut}(\mathbb{C}^3)_\omega$. Set $\eta = d\omega = dz_0 \wedge dz_1$. Remark that the invariance of ω implies the invariance of η and as a consequence the equality $(\phi_0, \phi_1)^* \eta = \eta$.

Proposition 3.1. *If ϕ belongs to $\text{Aut}(\mathbb{C}^3)_\omega$, then $\phi_* \frac{\partial}{\partial z_2} = \frac{\partial}{\partial z_2}$.*

In particular if ϕ belongs to $\text{Aut}(\mathbb{C}^3)_\omega$, then

$$\phi = (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1))$$

and the map

$$\varsigma: \text{Aut}(\mathbb{C}^3)_\omega \longrightarrow \text{Aut}(\mathbb{C}^2)_\eta, \quad (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1)) \mapsto (\phi_0(z_0, z_1), \phi_1(z_0, z_1))$$

is a morphism.

Proof. As we already mentioned, for a contact form there exists a unique vector field χ , called Reeb vector field, such that $\omega(\chi) = 1$ and $i_\chi d\omega = 0$; here $\chi = \frac{\partial}{\partial z_2}$. If ϕ belongs to $\text{Aut}(\mathbb{C}^3)_\omega$, then $\phi_*\chi = \chi$. As a result ϕ has the following form

$$\phi = (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1))$$

with (ϕ_0, ϕ_1) in $\text{Aut}(\mathbb{C}^2)$ and b in $\mathbb{C}[z_0, z_1]$. $\square$

Remark 3.2. Any element of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ can be written

$$(\varphi_0, \varphi_1, \det \text{jac } \varphi z_2 + b(z_0, z_1))$$

where $\varphi = (\varphi_0, \varphi_1) \in \text{Aut}(\mathbb{C}^2)$ and $db = (\det \text{jac } \varphi) z_0 dz_1 - \varphi_0 d\varphi_1$. Let us still denote by ς the natural projection

$$\varsigma: \text{Aut}(\mathbb{C}^3)_{c(\omega)} \rightarrow \text{Aut}(\mathbb{C}^2).$$

An element ϕ of $\text{Bir}(\mathbb{P}^2)_\eta$ is **exact** if it can be lifted via ς to $\text{Bir}(\mathbb{P}^3)_\omega$, or equivalently if it belongs to $\text{im } \varsigma$.

Contrary to the birational case (Theorem 8.1) any element of $\text{Aut}(\mathbb{C}^2)$ can be lifted via ς to $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$. Since b is defined up to a constant we do not speak about the ς -lift but a ς -lift.

The following obvious statement describes the group $\text{Aut}(\mathbb{C}^3)_\omega$:

Proposition 3.3. *Let us consider the morphism*

$$\varsigma: \text{Aut}(\mathbb{C}^3)_\omega \longrightarrow \text{Aut}(\mathbb{C}^2)_\eta, \quad (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1)) \mapsto (\phi_0(z_0, z_1), \phi_1(z_0, z_1)).$$

One has the following exact sequence

$$0 \longrightarrow \mathbb{C} \longrightarrow \text{Aut}(\mathbb{C}^3)_\omega \xrightarrow{\varsigma} \text{Aut}(\mathbb{C}^2)_\eta \longrightarrow 1; \quad (3.1)$$

more precisely $\ker \varsigma = \{(z_0, z_1, z_2 + \beta) \mid \beta \in \mathbb{C}\}$. In particular

$$\text{Aut}(\mathbb{C}^3)_\omega \simeq \text{Aut}(\mathbb{C}^2)_\eta \ltimes \mathbb{C}.$$

Proof. The 1-form $\phi_0 d\phi_1 - z_0 dz_1$ is a closed and polynomial one, so it is exact. Therefore ς is surjective. $\square$

Let G be a group. The **derived group** of G is the subgroup of G generated by all the commutators of G :

$$[G, G] = \langle ghg^{-1}h^{-1} \mid g, h \in G \rangle$$

The group G is said to be **perfect** if it coincides with its derived group, or equivalently, if the group has no nontrivial abelian quotients.

Such a property was established in the context of real smooth manifolds: Banyaga proved that the derived group of the group of contact diffeomorphisms is a perfect one ([2, 3, 4]).

Theorem 3.4. *The group $[\text{Aut}(\mathbb{C}^3)_\omega, \text{Aut}(\mathbb{C}^3)_\omega]$ is perfect.*

Proof. Since ς is surjective (Proposition 3.3) and $\text{Aut}(\mathbb{C}^2)_\eta$ is perfect ([21, Proposition 10]) the restriction of ς

$$\tilde{\varsigma} = \varsigma|_{[\text{Aut}(\mathbb{C}^3)_\omega, \text{Aut}(\mathbb{C}^3)_\omega]} : [\text{Aut}(\mathbb{C}^3)_\omega, \text{Aut}(\mathbb{C}^3)_\omega] \longrightarrow \text{Aut}(\mathbb{C}^2)_\eta$$

is surjective. Let ϕ be in $\ker \tilde{\varsigma}$; on the one hand $\phi = (z_0, z_1, z_2 + \beta)$ for some β (Proposition 3.3), and on the other hand ϕ is a product of commutators hence $\beta = 0$. We thus have the following exact sequence

$$0 \longrightarrow [\text{Aut}(\mathbb{C}^3)_\omega, \text{Aut}(\mathbb{C}^3)_\omega] \longrightarrow \text{Aut}(\mathbb{C}^2)_\eta \longrightarrow 1$$

and $[\text{Aut}(\mathbb{C}^3)_{\omega}, \text{Aut}(\mathbb{C}^3)_{\omega}] \simeq \text{Aut}(\mathbb{C}^2)_{\eta}$ which is perfect ([21, Proposition 10]). $\square$

3.2. Description of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$. Let us recall that $\text{Aut}(\mathbb{C}^2)$ is generated by J_2 and Aff_2 (see [27]). This implies that Aff_2 and

$$[J_2, J_2] = \{(z_0 + \beta, z_1 + P(z_0)) \mid \beta \in \mathbb{C}, P \in \mathbb{C}[z_0]\}.$$

generate $\text{Aut}(\mathbb{C}^2)$.

Proposition 3.5. *The group $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ is generated by $\mathcal{A}$ and $\mathcal{E}$ where*

$$\mathcal{E} = \{\zeta\text{-lifts of } \epsilon \mid \epsilon \in [J_2, J_2]\} \quad \text{and} \quad \mathcal{A} = \{\zeta\text{-lifts of } \alpha \mid \alpha \in \text{Aff}_2\}.$$

Proof. Let φ be a polynomial automorphism of $\mathbb{C}^2$ and let ϕ be a ζ -lift of φ to $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$

$$\phi = (\varphi, \det \text{jac } \varphi z_2 + b(z_0, z_1))$$

with b in $\mathbb{C}[z_0, z_1]$. One can write φ as $\alpha_1 \epsilon_1 \alpha_2 \epsilon_2 \dots \alpha_s \epsilon_s$ where α_i belongs to Aff_2 and ϵ_i to $[J_2, J_2]$. Let us now consider A_i a ζ -lift of α_i , $E_i = (\epsilon_i, z_2 + d_i)$ a ζ -lift of ϵ_i . Then $A_1 E_1 A_2 E_2 \dots A_s E_s$ belongs to $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$, and up to composition by an element $(z_0, z_1, z_2 + \beta) \in \mathcal{A}$ one has

$$\phi = A_1 E_1 A_2 E_2 \dots A_s E_s.$$

$\square$

Proposition 3.6. *One has*

$$\text{Aut}(\mathbb{C}^3)_{c(\omega)} \simeq \text{Aut}(\mathbb{C}^3)_{\omega} \ltimes \mathbb{C}^*.$$

Proof. Let us consider an element ϕ of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$, then $\phi^* \omega = V(\phi) \omega$ for some polynomial $V(\phi)$. As ω and $\phi^* \omega$ do not vanish, $V(\phi)$ does not vanish; therefore $V(\phi) = \lambda \in \mathbb{C}^*$. Let us write ϕ as follows:

$$\phi = (\lambda z_0, z_1, \lambda z_2) \circ \tilde{\phi};$$

of course $\tilde{\phi}^* \omega = \omega$. $\square$

Theorem 3.7. *The derived group $[\text{Aut}(\mathbb{C}^3)_{c(\omega)}, \text{Aut}(\mathbb{C}^3)_{c(\omega)}]$ of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ is perfect.*

Proof. According to Proposition 3.6 an element ϕ of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ can be written

$$(\lambda \phi_0, \phi_1, \lambda z_2 + \lambda b)$$

with $\lambda \in \mathbb{C}^*$ and $(\phi_0, \phi_1, z_2 + b) \in \text{Aut}(\mathbb{C}^3)_{\omega}$. Denote by φ the element of $\text{Aut}(\mathbb{C}^2)$ given by (ϕ_0, ϕ_1) . If ϕ belongs to $\ker \zeta$, then $\lambda = 1$, $\varphi = \text{id}$ and $b \in \mathbb{C}$, that is $\ker \zeta \simeq \mathbb{C}$ and

$$\mathbb{C} \longrightarrow \text{Aut}(\mathbb{C}^3)_{c(\omega)} \xrightarrow{\zeta} \text{Aut}(\mathbb{C}^2) \longrightarrow 1. \quad (3.2)$$

Since $\text{Aut}(\mathbb{C}^2)_{\eta}$ is perfect the restriction of ζ to $[\text{Aut}(\mathbb{C}^3)_{c(\omega)}, \text{Aut}(\mathbb{C}^3)_{c(\omega)}]$ induces the following exact sequence

$$0 \longrightarrow [\text{Aut}(\mathbb{C}^3)_{c(\omega)}, \text{Aut}(\mathbb{C}^3)_{c(\omega)}] \longrightarrow \text{Aut}(\mathbb{C}^2)_{\eta} \longrightarrow 1$$

and $[\text{Aut}(\mathbb{C}^3)_{c(\omega)}, \text{Aut}(\mathbb{C}^3)_{c(\omega)}] \simeq \text{Aut}(\mathbb{C}^2)_{\eta}$. One concludes as previously with [21, Proposition 10]. $\square$

3.3. Finite subgroups.

Proposition 3.8. *Any element of $\text{Aut}(\mathbb{C}^2)_\eta$ of period ℓ lifts via ς to a unique element of $\text{Aut}(\mathbb{C}^3)_\omega$ of period ℓ .*

Proof. Let us consider an element $\varphi = (\phi_0(z_0, z_1), \phi_1(z_0, z_1))$ of $\text{Aut}(\mathbb{C}^2)_\eta$. According to Theorem 8.1 there exists $b \in \mathbb{C}(z_0, z_1)$ such that $(\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1) + \mu)$ belongs to $\text{Bir}(\mathbb{P}^3)_\omega$ for any $\mu \in \mathbb{C}$. Assume that φ is of prime order ℓ ; let us prove that there exists a unique $\gamma \in \mathbb{C}$ such that

$$(\phi_0, \phi_1, z_2 + b(z_0, z_1) + \gamma)$$

is of order ℓ .

Assume for simplicity that $\ell = 2$ (but a similar argument works for any ℓ). Let us recall that the following equality holds

$$z_0 dz_1 - \phi_0 d\phi_1 = db \quad (3.3)$$

Applying ϕ to this equality one gets

$$\phi_0 d\phi_1 - z_0 dz_1 = d(b \circ \varphi) \quad (3.4)$$

We add (3.3) and (3.4) and obtain that $b + b \circ \varphi$ is a constant β . Furthermore

$$(\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1) + \mu)^2 = (z_0, z_1, z_2 + 2\gamma + b + b \circ \varphi) = (z_0, z_1, z_2 + 2\gamma + \beta)$$

so as soon as $\gamma = -\beta/2$ one has $(\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1) + \mu)^2 = \text{id}$. $\square$

Proposition 3.9. *A finite subgroup of $\text{Aut}(\mathbb{C}^2)$ can be lifted to a finite subgroup of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$.*

Proof. Let H be a finite subgroup of $\text{Aut}(\mathbb{C}^2)$. The group H is linearizable ([24]) hence has a fixed point p . Since the translations belong to $\text{Aut}(\mathbb{C}^2)$ one can assume that $p = (0, 0)$. Let us consider the lifts of all elements of H in $(\text{Aut}(\mathbb{C}^3)_{c(\omega)}|_0)$; they form a group isomorphic to H so is in particular finite. $\square$

Remark 3.10. Any subgroup G of $\text{Aut}(\mathbb{C}^2)$ that preserves $(0, 0)$ can be lifted to a subgroup of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ isomorphic to G .

Theorem 3.11. *Any finite subgroup of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ is linearizable via an element of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$.*

Proof. Let G be a finite subgroup of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$. The group G is isomorphic to $H = \varsigma(G)$ which is thus a finite subgroup of $\text{Aut}(\mathbb{C}^2)$. There exists a map $h \in \text{Aut}(\mathbb{C}^2)$ that linearizes H (see [24]); as a result H has a fixed point p and up to translations one can suppose that $p = (0, 0)$. Note that $h(0) = 0$. The lift of h in $(\text{Aut}(\mathbb{C}^3)_{c(\omega)}|_0)$ linearizes G . $\square$

4. AUTOMORPHISMS GROUP

Let us first introduce some notations. The group of the field automorphisms of $\mathbb{C}$ acts on $\text{Aut}(\mathbb{C}^n)$ (resp. $\text{Bir}(\mathbb{P}^n)$): if f is an element of $\text{Aut}(\mathbb{C}^n)$ and if ξ is a field automorphism we denote by ${}^\xi f$ the element obtained by letting ξ acting on f . Using the structure of amalgamated product of $\text{Aut}(\mathbb{C}^2)$, the automorphisms of this group have been described ([16]): let φ be an automorphism of $\text{Aut}(\mathbb{C}^2)$; there exist a polynomial automorphism ψ of $\mathbb{C}^2$ and a field automorphism ξ such that

$$\forall f \in \text{Aut}(\mathbb{C}^2) \quad \varphi(f) = {}^\xi(\psi f \psi^{-1}).$$

Even if $\text{Bir}(\mathbb{P}^2)$ has not the same structure as $\text{Aut}(\mathbb{C}^2)$ (see Appendix of [12]) the automorphisms group of $\text{Bir}(\mathbb{P}^2)$ can be described and a similar result is obtained ([17]). Then, in 2011, Kraft and Stampfli show that every automorphism of $\text{Aut}(\mathbb{C}^n)$ is inner up to field automorphisms when restricted to the group Tame_n (see [28]).

We now would like to describe the group $\text{Aut}(\text{Aut}(\mathbb{C}^3)_{\omega})$. Let us recall that the **center** of a group G , denoted $Z(G)$, is the set of elements that commute with every element of G .

Proposition 4.1. *The center of $\text{Aut}(\mathbb{C}^3)_{\omega}$ is isomorphic to $\mathbb{C}$:*

$$Z(\text{Aut}(\mathbb{C}^3)_{\omega}) = \{(z_0, z_1, z_2 + \beta) \mid \beta \in \mathbb{C}\}$$

and the center of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ is trivial.

As $\text{Aut}(\mathbb{C}^3)_{\omega} \simeq \text{Aut}(\mathbb{C}^2)_{\eta} \ltimes \mathbb{C}$ Proposition 4.1 implies the following statement:

Corollary 4.2. *The quotient of $\text{Aut}(\mathbb{C}^3)_{\omega}$ by its center is isomorphic to $\text{Aut}(\mathbb{C}^2)_{\eta}$.*

Lemma 4.3. *One has the following isomorphism*

$$\text{Hom}(\text{Aut}(\mathbb{C}^3)_{\omega}, \mathbb{C}) \simeq \text{Hom}(\mathbb{C}, \mathbb{C})$$

where $\text{Hom}(\mathbb{C}, \mathbb{C})$ denotes the homomorphisms of the additive group $\mathbb{C}$.

Proof. Note that if ϕ belongs to $[\text{Aut}(\mathbb{C}^3)_{\omega}, \text{Aut}(\mathbb{C}^3)_{\omega}]$, then the last component of ϕ is well defined (that is not defined modulo a constant). Besides $\text{Aut}(\mathbb{C}^3)_{\omega} \simeq \text{Aut}(\mathbb{C}^2)_{\eta} \ltimes \mathbb{C}$ and $\text{Aut}(\mathbb{C}^2)_{\eta}$ is perfect thus

$$\text{Aut}(\mathbb{C}^3)_{\omega} / [\text{Aut}(\mathbb{C}^3)_{\omega}, \text{Aut}(\mathbb{C}^3)_{\omega}] \simeq \mathbb{C}$$

and

$$\begin{array}{ccc} \text{Aut}(\mathbb{C}^3)_{\omega} \simeq \text{Aut}(\mathbb{C}^2)_{\eta} \ltimes \mathbb{C} & & \\ \downarrow & \searrow & \\ \text{Aut}(\mathbb{C}^3)_{\omega} / [\text{Aut}(\mathbb{C}^3)_{\omega}, \text{Aut}(\mathbb{C}^3)_{\omega}] & \xrightarrow{\sim} & \mathbb{C} \end{array}$$

We conclude by noting that any element of $\text{Hom}(\text{Aut}(\mathbb{C}^3)_{\omega}, \mathbb{C})$ acts trivially on ϕ . □

Remark 4.4. An element c of $\text{Hom}(\text{Aut}(\mathbb{C}^3)_{\omega}, \mathbb{C})$ acts on $\text{Aut}(\mathbb{C}^3)_{\omega}$ as follows

$$(\phi_0, \phi_1, z_3 + b(z_0, z_1)) \rightarrow (\phi_0, \phi_1, z_3 + b(z_0, z_1) + c(\phi))$$

Definition. Let H be a normal subgroup of a group G . We say that an automorphism of H of the form $\phi \mapsto \varphi\phi\varphi^{-1}$, with φ in G , is **G -inner**.

Theorem 4.5. *The group $\text{Aut}(\text{Aut}(\mathbb{C}^3)_{\omega})$ is generated by the automorphisms group of the field $\mathbb{C}$, the group of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ -inner automorphisms and the action of $\text{Hom}(\mathbb{C}, \mathbb{C})$.*

Proof. Consider an element ψ of $\text{Aut}(\text{Aut}(\mathbb{C}^3)_{\omega})$. For any $\phi = (\varphi_{\phi}, z_2 + T_{\phi}(z_0, z_1))$ one has

$$\psi(\phi) = (\widetilde{\varphi}_{\phi}, z_2 + \Delta_{\phi}(z_0, z_1)).$$

In particular ψ induces an automorphism ψ_0 of $\text{Aut}(\mathbb{C}^2)_{\eta}$; indeed since ψ is an automorphism of $\text{Aut}(\mathbb{C}^3)_{\omega}$, it preserves $Z(\text{Aut}(\mathbb{C}^3)_{\omega})$ and so, from Corollary 4.2 induces an automorphism of $\text{Aut}(\mathbb{C}^2)_{\eta}$.

According to Theorem 13.2 one can assume that $\psi_0 = \text{id}$ up to the action of an automorphism of the field $\mathbb{C}$ and up to conjugacy by an $\text{Aut}(\mathbb{C}^2)$ -inner automorphism, i.e.

$$\psi(\phi) = (\varphi_{\phi}, z_2 + \Delta_{\phi}(z_0, z_1))$$

Set $\phi^{-1} = (\phi_0^{-1}, z_2 + T_{\phi^{-1}}(z_0, z_1))$. On the one hand $\phi^{-1} \circ \phi = (\text{id}, z_2 + T_{\phi}(z_0, z_1) + T_{\phi^{-1}}(\phi_0))$ so

$$T_{\phi} + T_{\phi^{-1}}(\phi_0) = 0 \quad (4.1)$$

and on the other hand

$$\psi(\phi \circ \phi^{-1}) = (\text{id}, z_2 + T_{\phi^{-1}}(z_0, z_1) + \Delta_{\phi} \phi_0^{-1})$$

belongs to $\text{Aut}(\mathbb{C}^3)_{\omega}$ hence $T_{\phi^{-1}} + \Delta_{\phi} \phi_0^{-1}$ is a constant. This, combined with (4.1), implies that $\Delta_{\phi} = T_{\phi} + c_{\phi}$, where c_{ϕ} is a constant, and yields to a morphism from $\text{Aut}(\mathbb{C}^3)_{\omega}$ to $\mathbb{C}$:

$$\text{Aut}(\mathbb{C}^3)_{\omega} \rightarrow \mathbb{C}, \quad \phi \mapsto c_{\phi}.$$

Consider an homomorphism

$$\rho: \text{Aut}(\mathbb{C}^3)_{\omega} \rightarrow \mathbb{C}, \quad \phi \mapsto \rho_{\phi}.$$

Let us define $\psi: \text{Aut}(\mathbb{C}^3)_{\omega} \rightarrow \text{Aut}(\mathbb{C}^3)_{\omega}$ by:

$$\psi(\phi) = (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1) + \rho_{\phi})$$

where $\phi = (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1)) \in \text{Aut}(\mathbb{C}^3)_{\omega}$. One can check that ψ belongs to $\text{Aut}(\text{Aut}(\mathbb{C}^3)_{\omega})$. $\square$

Part 2. Contact birational maps

A **rational map** of $\mathbb{P}^n$ can be written

$$\phi: \mathbb{P}^n \dashrightarrow \mathbb{P}^n \quad (z_0 : z_1 : \dots : z_n) \dashrightarrow (\phi_0(z_0, z_1, \dots, z_n) : \phi_1(z_0, z_1, \dots, z_n) : \dots : \phi_n(z_0, z_1, \dots, z_n))$$

where the ϕ_i 's are homogeneous polynomials of the same degree ≥ 1 and without common factor of positive degree. The **degree** of ϕ is by definition the degree of the ϕ_i . A **birational map** of $\mathbb{P}^n$ is a rational map that admits a rational inverse. Of course $\text{Aut}(\mathbb{C}^n)$ is a subgroup of $\text{Bir}(\mathbb{P}^n)$. An other natural subgroup of $\text{Bir}(\mathbb{P}^n)$ is the group $\text{Aut}(\mathbb{P}^n) \simeq \text{PGL}(n+1; \mathbb{C})$ of automorphisms of $\mathbb{P}^n$.

The **indeterminacy set** $\text{Ind}\phi$ of ϕ is the set of the common zeros of the ϕ_i 's. The **exceptional set** $\text{Exc}\phi$ of ϕ is the (finite) union of subvarieties M_i of $\mathbb{P}^n$ such that ϕ is not injective on any open subset of M_i .

Let us extend the definition of Jonquières group we gave in the case of polynomial automorphisms of $\mathbb{C}^n$ to the case of birational maps of $\mathbb{P}^2$: the **Jonquières group**, denoted $\mathcal{J}$, is the group of birational maps of $\mathbb{P}^2$ that preserve a pencil of rational curves. Since two pencils of rational curves are birationally conjugate, $\mathcal{J}$ does not depend, up to conjugacy, of the choice of the pencil. In other words one can decide, up to birational conjugacy, that $\mathcal{J}$ is in the affine chart $z_2 = 1$ the maximal group of birational maps that preserve the fibration $z_1 = \text{cst}$. An element ϕ of $\mathcal{J}$ permutes the fibers of the fibration thus induces an automorphism of the base $\mathbb{P}^1$; note that if the fibration is fiberwise invariant, ϕ acts as an homography in the generic fibers. Hence $\mathcal{J}$ can be identified with the semi-direct product $\text{PGL}(2; \mathbb{C}(z_1)) \rtimes \text{PGL}(2; \mathbb{C})$.

We study the birational maps $\phi = (\phi_0, \phi_1, \phi_2)$ defined on $\mathbb{C}^3 = (z_3 = 1) \subset \mathbb{P}^3$ that preserve either the contact standard form ω , or the contact structure $c(\omega)$ associated to ω . In other words we would like to describe the groups $\text{Bir}(\mathbb{P}^3)_{\omega}$ and $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ and also their elements.

Let us now illustrate a fundamental difference between $\text{Bir}(\mathbb{P}^3)_{\omega}$ and $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$: the first group preserves the fibration associated to $\frac{\partial}{\partial z_2}$ whereas the second doesn't.

Proposition 4.6. *If ϕ belongs to $\text{Bir}(\mathbb{P}^3)_{\omega}$, then $\phi_* \frac{\partial}{\partial z_2} = \frac{\partial}{\partial z_2}$.*

In particular if ϕ belongs to $\text{Bir}(\mathbb{P}^3)_{\omega}$, then

$$\phi = (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1))$$

and the map

$$\varsigma: \text{Bir}(\mathbb{P}^3)_{\omega} \longrightarrow \text{Bir}(\mathbb{P}^2)_{\eta}, \quad (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1)) \mapsto (\phi_0(z_0, z_1), \phi_1(z_0, z_1))$$

is a morphism.

Remark 4.7. The proof is similar to the proof of Proposition 3.1.

Remark 4.8. The first assertion of Proposition 4.6 is not true for the group $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$; indeed let us consider the map ψ defined by

$$\psi = \left(\frac{z_0}{(1+z_2)^2}, z_1, \frac{z_2}{1+z_2} \right);$$

it belongs to $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ and does not preserve the fibration associated to the vector field $\frac{\partial}{\partial z_2}$.

5. A P.D.E. APPROACH

Let $\phi = (\phi_0, \phi_1, \phi_2)$ be in $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$; then $\phi^* \omega = V(\phi) \omega$ for some rational function $V(\phi)$. One inherits a map V from $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ into the set of rational functions in z_0, z_1 and z_2 . The equality $\phi^* \omega = V(\phi) \omega$ gives the following system $(\star)$ of P. D. E.:

$$\begin{cases} \phi_0 \frac{\partial \phi_1}{\partial z_0} + \frac{\partial \phi_2}{\partial z_0} = 0 & (\star_1) \\ \phi_0 \frac{\partial \phi_1}{\partial z_1} + \frac{\partial \phi_2}{\partial z_1} = V(\phi) z_0 & (\star_2) \\ \phi_0 \frac{\partial \phi_1}{\partial z_2} + \frac{\partial \phi_2}{\partial z_2} = V(\phi) & (\star_3) \end{cases}$$

Thanks to $(\star_2)$ and $(\star_3)$ one gets

$$\phi_0 \left(\frac{\partial \phi_1}{\partial z_1} - z_0 \frac{\partial \phi_1}{\partial z_2} \right) + \left(\frac{\partial \phi_2}{\partial z_1} - z_0 \frac{\partial \phi_2}{\partial z_2} \right) = 0 \quad (\star_4)$$

Equation $(\star_1)$ has a special family of solutions: maps for which both ϕ_1 or ϕ_2 do not depend on z_0 (note that if ϕ_1 (resp. ϕ_2) does not depend on z_0 then $(\star_1)$ implies that ϕ_2 (resp. ϕ_1) also; in that case we can then compute ϕ_0 thanks to $(\star_4)$). Taking (ϕ_1, ϕ_2) in $\text{Bir}(\mathbb{P}^2)$ we get elements in $\text{im } \mathcal{K}$; we will called this family of solutions **Klein family**. Note that this family is a group denoted $\mathcal{K}$, the **Klein group**.

Proposition 5.1. *The elements of $\mathcal{K}$ are of the following type*

$$\left(\frac{-\frac{\partial \phi_2}{\partial z_1} + z_0 \frac{\partial \phi_2}{\partial z_2}}{\frac{\partial \phi_1}{\partial z_1} - z_0 \frac{\partial \phi_1}{\partial z_2}}, \phi_1(z_1, z_2), \phi_2(z_1, z_2) \right)$$

with (ϕ_1, ϕ_2) in $\text{Bir}(\mathbb{P}^2)$.

Assume now that ϕ_1 or ϕ_2 really depends on z_0 (i.e. that ϕ does not belong to the Klein family). Then $(\star_1)$ and $(\star_4)$ imply

$$\left(\frac{\partial \phi_2}{\partial z_1} - z_0 \frac{\partial \phi_2}{\partial z_2} \right) \frac{\partial \phi_1}{\partial z_0} = \left(\frac{\partial \phi_1}{\partial z_1} - z_0 \frac{\partial \phi_1}{\partial z_2} \right) \frac{\partial \phi_2}{\partial z_0} \quad (\star_5)$$

One can rewrite $(\star_5)$ as

$$\frac{\frac{\partial \phi_2}{\partial z_1} - z_0 \frac{\partial \phi_2}{\partial z_2}}{\frac{\partial \phi_2}{\partial z_0}} = \frac{\frac{\partial \phi_1}{\partial z_1} - z_0 \frac{\partial \phi_1}{\partial z_2}}{\frac{\partial \phi_1}{\partial z_0}}.$$

Denote by α the map from $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ to the set of rational functions in z_0, z_1 and z_2 defined by $\alpha(\phi) = \infty$ if ϕ belongs to $\mathcal{K}$ and

$$\alpha(\phi) = \frac{\frac{\partial \phi_2}{\partial z_1} - z_0 \frac{\partial \phi_2}{\partial z_2}}{\frac{\partial \phi_2}{\partial z_0}} = \frac{\frac{\partial \phi_1}{\partial z_1} - z_0 \frac{\partial \phi_1}{\partial z_2}}{\frac{\partial \phi_1}{\partial z_0}}$$

otherwise.

If ϕ_1 and ϕ_2 are some first integrals of

$$Z_\phi = \alpha(\phi) \frac{\partial}{\partial z_0} - \frac{\partial}{\partial z_1} + z_0 \frac{\partial}{\partial z_2},$$

then $(\star_5)$ is satisfied. One thus gets ϕ_0 from $(\star_1)$. Note that such a ϕ is not always birational. But one can get a lot of birational examples by this way.

For instance when $\alpha(\phi) \equiv 0$ one obtains a family of rational maps solutions of $(\star)$ and Legendre involution is one of them. The set of birational maps of that family is called **Legendre family**, i.e. it is the set of birational maps of the following form

$$\left(-\frac{\frac{\partial}{\partial z_0}(\phi_2(z_0, -(z_2 + z_0 z_1)))}{\frac{\partial}{\partial z_0}(\phi_1(z_0, -(z_2 + z_0 z_1)))}, \phi_1(z_0, -(z_2 + z_0 z_1)), \phi_2(z_0, -(z_2 + z_0 z_1)) \right).$$

Remark 5.2. The Legendre family composed with the Legendre involution (right composition) yields to the Klein family.

Definition. Let γ be an irreducible curve; γ is a **legendrian curve** if $s_\gamma^* \omega = 0$ where s_γ denotes a local parametrization of γ .

Remark 5.3. Elements of the Klein family preserve the fibration $\{z_1 = \text{cst}, z_2 = \text{cst}\}$; note that its fibers are legendrian curves. The Legendre involution sends the fibration $\{z_0 = \text{cst}, z_2 + z_0 z_1 = \text{cst}\}$ onto $\{z_1 = \text{cst}, z_2 = \text{cst}\}$. Then of course if one conjugates the Klein family by the Legendre involution one gets a family that preserves the fibration by legendrian curves $\{z_0 = \text{cst}, z_2 + z_0 z_1 = \text{cst}\}$.

A direct computation implies:

Proposition 5.4. Let $\phi = (\phi_0, \phi_1, \phi_2)$ be a contact birational map of $\mathbb{P}^3$.

The map ϕ conjugates the foliation induced by Z_ϕ to the foliation induced by $\frac{\partial}{\partial z_0}$.

As a consequence the field of the rational first integrals of Z_ϕ is generated by ϕ_1 and ϕ_2 .

5.1. Actions of $\mathcal{K}$ and $\text{Bir}(\mathbb{P}^3)_\omega$ on $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$. The left translation action of $\mathcal{K}$ on $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ is given by

$$(\psi, \phi) \in \mathcal{K} \times \text{Bir}(\mathbb{P}^3)_{c(\omega)} \longrightarrow \psi\phi \in \text{Bir}(\mathbb{P}^3)_{c(\omega)}.$$

Take ϕ and ψ in $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ such that $\alpha(\phi) = \alpha(\psi)$, then ψ_1 and ψ_2 are first integrals of Z_ϕ and by Proposition 5.4

$$\psi_1 = \phi_1(\phi_1, \phi_2), \quad \psi_2 = \phi_2(\phi_1, \phi_2)$$

where $\phi = (\phi_1, \phi_2)$ is birational. Hence

$$\psi\phi^{-1} = (\psi_0 \circ \phi^{-1}, \phi_1(z_1, z_2), \phi_2(z_1, z_2))$$

belongs to $\mathcal{K}$; in other words ϕ and ψ are in the same $\mathcal{K}$ -orbit.

Assume now that $\psi = \kappa\phi$ where κ denotes an element of $\mathcal{K}$. Then the foliations defined by Z_ϕ and Z_ψ coincide because they have the same set of first integrals. As a consequence $\alpha(\phi) = \alpha(\psi)$.

Hence one can state:

Theorem 5.5. *The map α is a complete invariant of the left translation action of $\mathcal{K}$ on $\text{Bir}(\mathbb{P}^3)_{\mathbb{C}(\omega)}$, that is for any ϕ and ψ in $\text{Bir}(\mathbb{P}^3)_{\mathbb{C}(\omega)}$ one has $\alpha(\phi) = \alpha(\psi)$ if and only if $\psi\phi^{-1}$ belongs to $\mathcal{K}$.*

Question 1. *Is the map α surjective ?*

Let us consider the following differential equation

$$y'' = F(x, y, y') \quad (5.1)$$

where F denotes a rational function. Set $y' = u$, then

$$(5.1) \Leftrightarrow \begin{cases} \frac{du}{dt} = F(x, y, u) \\ \frac{dy}{dt} = u \\ \frac{dx}{dt} = 1 \end{cases}$$

So one can associate to (5.1) the following vector field

$$Z = F \frac{\partial}{\partial u} + u \frac{\partial}{\partial y} + \frac{\partial}{\partial x}.$$

We say that (5.1) is **rationally integrable** if the vector field Z has two first integrals r_1 and r_2 rationally independent: $dr_1 \wedge dr_2 \neq 0$.

For generic γ and β in $\mathbb{C}$ the differential equation $y'' + \gamma y' + \beta y = 0$ is not rationally integrable; as a consequence $-\gamma z_0 - \beta z_2$ is not in the image of α . The first Painlevé equation gives examples of polynomial of degree 2 that does not belong to $\text{im } \alpha$:

Theorem 5.6 ([?]). *The equation $\mathcal{P}_1$*

$$y'' = 6y^2 + x$$

is not rationally integrable.

If we come back with our notations it means that $6z_2^2 - z_1$ is not in the image of α .

Remark 5.7. Indeed all generic Painlevé equations give rise to rational functions that do not belong to $\text{im } \alpha$.

Nevertheless one can easily obtain examples of elements in the image of α :

Examples 5.8. – If $\phi = \left(\frac{z_0}{\beta}, z_0 + \beta z_1, z_2 - \frac{z_0^2}{2\beta} \right)$ with $\beta \in \mathbb{C}^*$, then $\alpha(\phi) = \beta$.

– If

$$\phi = (z_0, z_1 + P(z_0), z_2 + Q(z_0))$$

with P, Q in $\mathbb{C}[z_0]$ such that $Q'(z_0) = -z_0 P'(z_0)$, then $\alpha(\phi) = \frac{1}{P'(z_0)}$.

– If

$$\phi = (-z_1, z_0 + P(z_1), z_2 + z_0 z_1 + Q(z_1))$$

with P, Q in $\mathbb{C}[z_1]$ such that $Q'(z_1) = z_1 P'(z_1)$ then $\alpha(\phi) = P'(z_1)$.

Consider the left translation action of $\text{Bir}(\mathbb{P}^3)_{\omega}$ on $\text{Bir}(\mathbb{P}^3)_{\mathbb{C}(\omega)}$ defined by

$$(\psi, \phi) \in \text{Bir}(\mathbb{P}^3)_{\omega} \times \text{Bir}(\mathbb{P}^3)_{\mathbb{C}(\omega)} \longrightarrow \psi\phi \in \text{Bir}(\mathbb{P}^3)_{\mathbb{C}(\omega)}.$$

Theorem 5.9. *The map V is a complete invariant of the left translation action of $\text{Bir}(\mathbb{P}^3)_{\omega}$ on $\text{Bir}(\mathbb{P}^3)_{\mathbb{C}(\omega)}$: for any ϕ, ψ in $\text{Bir}(\mathbb{P}^3)_{\mathbb{C}(\omega)}$ one has $V(\phi) = V(\psi)$ if and only if $\psi\phi^{-1}$ belongs to $\text{Bir}(\mathbb{P}^3)_{\omega}$.*

Proof. Let ϕ be a contact birational map of $\mathbb{P}^3$. Obviously $(f\phi)^*\omega = V(\phi)\omega$ for any $f \in \text{Bir}(\mathbb{P}^3)_\omega$.

Let us now consider two contact birational maps ϕ and ψ of $\mathbb{P}^3$ such that $V = V(\phi) = V(\psi)$. On the one hand

$$(\phi^{-1})^*\psi^*\omega = (\phi^{-1})^*V(\phi)\omega = V \circ \phi^{-1}(\phi^{-1})^*\omega$$

and on the other hand composing $\phi^*\omega = V\omega$ by $(\phi^{-1})^*$ one gets

$$\phi^*\omega = V\omega \Rightarrow (\phi^{-1})^*(\phi^*\omega) = (\phi^{-1})^*(V\omega) \Rightarrow \omega = V \circ \phi^{-1}(\phi^{-1})^*\omega.$$

As a consequence $(\phi^{-1})^*\psi^*\omega = \omega$, that is $\psi\phi^{-1}$ belongs to $\text{Bir}(\mathbb{P}^3)_\omega$. $\square$

Proposition 5.10. *If ϕ and ψ are two contact birational maps of $\mathbb{P}^3$ such that $\alpha(\phi) = \alpha(\psi)$ and $V(\phi) = V(\psi)$, then $\psi\phi^{-1}$ belongs to*

$$\left\{ \left(\frac{z_0 - b'(z_1)}{v'(z_1)}, v(z_1), z_2 + b(z_1) \right) \mid b \in \mathbb{C}(z_1), v \in \text{PGL}(2; \mathbb{C}) \right\} = \mathcal{K} \cap \text{Bir}(\mathbb{P}^3)_\omega.$$

Proof. Since both $\alpha(\phi) = \alpha(\psi)$ and $V(\phi) = V(\psi)$ the map $\psi\phi^{-1}$ is an element of $\text{Bir}(\mathbb{P}^3)_\omega \cap \mathcal{K}$. One gets the result from the descriptions of the Klein family and of $\text{Bir}(\mathbb{P}^3)_\omega$ (Proposition 3.1). $\square$

Let us now give some examples of $V(\phi)$.

Examples 5.11. – If ϕ belongs to $\mathcal{K}$, then

$$V(\phi) = \frac{\frac{\partial\phi_1}{\partial z_1} \frac{\partial\phi_2}{\partial z_2} - \frac{\partial\phi_1}{\partial z_2} \frac{\partial\phi_2}{\partial z_1}}{\frac{\partial\phi_1}{\partial z_1} - z_0 \frac{\partial\phi_1}{\partial z_2}}.$$

– If

$$\phi = \left(\frac{1}{nz_0^{n-1}z_2 + (n+1)z_0^n(z_1+1)}, z_0^n(z_0 + z_2 + z_0z_1), -z_0 \right)$$

$$\text{with } n \in \mathbb{Z}, \text{ then } V(\phi) = \frac{z_0}{(n+1)z_0z_1 + nz_2 + (n+1)z_0}.$$

– If

$$\phi = \left(\frac{(z_1 - z_0)^2}{2z_0z_1 + 2z_2 - z_0^2}, \frac{2z_2 + z_0^2}{z_1 - z_0}, z_1 - z_0 \right),$$

$$\text{then } V(\phi) = \frac{2(z_0 - z_1)}{z_0^2 - 2z_0z_1 - 2z_2}.$$

Remark 5.12. If ϕ belongs to $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$, then $\phi^*\omega = V(\phi)\omega$ and $\phi^*(\omega \wedge d\omega) = V(\phi)^2\omega \wedge d\omega$ and $\det \text{jac } \phi$ is a square. This gives some constraint on $V(\phi)$.

As previously we can ask: is V surjective? The answer is no. Indeed let us assume that there exists $\phi \in \text{Bir}(\mathbb{P}^3)_{c(\omega)}$ such that $V(\phi) = z_2$. Then $\phi_0 d\phi_0 + d\phi_2 = z_0z_2 dz_1 + d\left(\frac{z_2^2}{2}\right)$ and $d\phi_0 \wedge d\phi_1 = d(z_0z_2) \wedge dz_1$. Since the fibers of (z_0z_2, z_1) are connected one can write ϕ_0 as $\phi_0(z_0z_2, z_1)$ and ϕ_1 as $\phi_1(z_0z_2, z_1)$. Then $\phi^*\omega = V(\phi)\omega$ implies that $\phi_2 - \frac{z_2^2}{2} = \phi_2(z_0z_2, z_1)$. In other words

$$\phi = \left(\phi_0(z_0z_2, z_1), \phi_1(z_0z_2, z_1), \phi_2(z_0z_2, z_1) + \frac{z_2^2}{2} \right).$$

But $\phi \circ \left(\frac{z_0}{z_2}, z_1, z_2 \right)$ is clearly not birational so does ϕ : contradiction.

6. INVARIANT FORMS AND VECTOR FIELDS

The next statement deals with flows in $\text{Bir}(\mathbb{P}^3)_{\omega}$ (see [13] for a definition).

Proposition 6.1. *Let ϕ_t be a flow in $\text{Bir}(\mathbb{P}^3)_{\omega}$. Then ϕ_t has a first integral depending only on (z_0, z_1) and with rational fibers.*

In other words

$$\phi_t = (\phi_t(z_0, z_1), z_2 + b_t(z_0, z_1))$$

where ϕ_t belongs, up to conjugacy, to $\mathcal{J}$ and b_t to $\mathbb{C}(z_0, z_1)$.

Proof. Let χ be the infinitesimal generator of ϕ_t , i.e.

$$\chi = \left. \frac{\partial \phi_t}{\partial t} \right|_{t=0}.$$

By derivating $\phi_t^* \omega = \omega$ with respect to t one gets that the Lie derivative $L_{\chi} \omega$ is zero. Set $\chi = \sum_{i=0}^2 \chi_i \frac{\partial}{\partial z_i}$, hence

$$L_{\chi} \omega = \iota_{\chi} d\omega + d\iota_{\chi} \omega = \chi_0 dz_1 + z_0 d\chi_1 + d\chi_2$$

and so

$$L_{\chi} \omega = \left(z_0 \frac{\partial \chi_1}{\partial z_0} + \frac{\partial \chi_2}{\partial z_0} \right) dz_0 + \left(\chi_0 + z_0 \frac{\partial \chi_1}{\partial z_1} + \frac{\partial \chi_2}{\partial z_1} \right) dz_1 + \left(z_0 \frac{\partial \chi_1}{\partial z_2} + \frac{\partial \chi_2}{\partial z_2} \right) dz_2.$$

In particular $z_0 \chi_1 + \chi_2 = \gamma(z_0, z_1)$, then $\chi_0 + \frac{\partial}{\partial z_1}(z_0 \chi_1 + \chi_2) = 0$ so $\chi_0 = -\frac{\partial \gamma}{\partial z_1}$ and finally $\chi_1 = \frac{\partial \gamma}{\partial z_0}$.

If γ is constant, then $\chi = \frac{\partial}{\partial z_2}$, that is $\phi_t = (z_0, z_1, z_2 + \beta t)$ with $\beta \in \mathbb{C}$.

Let us now assume that γ is non-constant; one has

$$\chi = -\frac{\partial \gamma}{\partial z_1} \frac{\partial}{\partial z_0} + \frac{\partial \gamma}{\partial z_0} \frac{\partial}{\partial z_1} + \left(\gamma(z_0, z_1) - z_0 \frac{\partial \gamma}{\partial z_0} \right) \frac{\partial}{\partial z_2}$$

and γ is a first integral of χ . For all t

$$\phi_t = (\phi_{0,t}(z_0, z_1), \phi_{1,t}(z_0, z_1), z_2 + b_t(z_0, z_1))$$

and the function γ is invariant by ϕ_t and as a consequence by the flow ϕ_t . The fibers of γ in $\mathbb{C}^2$ (up to compactification/normalization) are rational or elliptic since they own a flow. As $\langle \phi_t \rangle$ is uncountable they have to be rational ([10]) and up to conjugacy ϕ_t belongs to $\mathcal{J}$. $\square$

The following examples contain many flows.

Example 6.2. The elements of $\text{Aut}(\mathbb{P}^3)_{c(\omega)}$ can be written

$$(\varepsilon z_0 + \lambda, \beta z_1 + \gamma, -\beta \lambda z_1 + \varepsilon \beta z_2 + \delta)$$

with ε, β in $\mathbb{C}^*$ and λ, γ, δ in $\mathbb{C}$. The group $\text{Aut}(\mathbb{P}^3)_{c(\omega)}$ acts transitively on $\mathbb{C}^3 = \{z_3 = 1\}$.

Examples 6.3. a) For any $\varepsilon, \beta, \gamma$ and δ in $\mathbb{C}$ such that $\varepsilon \delta - \beta \gamma \neq 0$, the map

$$\left(\frac{(\gamma z_1 + \delta)^2}{\varepsilon \delta - \beta \gamma} z_0, \frac{\varepsilon z_1 + \beta}{\gamma z_1 + \delta}, z_2 \right)$$

belongs to $\text{Bir}(\mathbb{P}^3)_{\omega}$. These maps form a group contained in $\text{im } \mathcal{K}$ and isomorphic to $\text{PGL}(2; \mathbb{C})$.

b) The birational maps given by

- $(z_0, z_1 + \phi(z_0), z_2 + \psi(z_0))$ with $z_0 \phi'(z_0) + \psi'(z_0) = 0$,
- $(z_0 - \psi'(z_1), z_1, z_2 + \psi(z_1))$

belong to $\text{Bir}(\mathbb{P}^3)_{\omega}$. Any of these families forms an abelian group.

The fact that an element of $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ preserves a vector field and the fact that it preserves a contact form are related:

Proposition 6.4. *Let ϕ be a contact birational map of $\mathbb{P}^3$. There exist a contact form Θ colinear to ω such that $\phi^*\Theta = \Theta$ if and only if $V(\phi)$ can be written $\frac{U}{U \circ \phi}$ for some rational function U . In that case ϕ preserves the Reeb flow associated to Θ , so a foliation by curves.*

Proof. Assume that such a Θ exists. On the one hand $\phi^*\omega = V(\phi)\omega$ and on the other hand $\Theta = U\omega$. Hence

$$\phi^*\Theta = U \circ \phi \cdot \phi^*\omega = U \circ \phi \cdot V(\phi)\omega = \frac{U \circ \phi}{U} \cdot V(\phi)\Theta$$

and so if such U exists, one has $V(\phi) = \frac{U}{U \circ \phi}$.

Reciprocally if $\phi \in \text{Bir}(\mathbb{P}^3)_{c(\omega)} \setminus \text{Bir}(\mathbb{P}^3)_\omega$ satisfies $\phi^*\omega = \frac{U}{U \circ \phi}\omega$ for some rational function U , then $\phi^*\Theta = \Theta$ where $\Theta = U\omega$. $\square$

Examples 6.5. – First consider the Legendre involution $\mathcal{L} = (z_1, z_0, -z_2 - z_0z_1)$. As we have seen $V(\mathcal{L}) = -1$. One can check that $U = z_2 + \frac{z_0z_1}{2}$ suits.

– For an element ϕ in $\text{Aut}(\mathbb{P}^3)_{c(\omega)}$

$$\phi = (\varepsilon z_0 + \lambda, \beta z_1 + \gamma, -\beta \lambda z_1 + \varepsilon \beta z_2 + \delta)$$

with ε, β in $\mathbb{C}^*$ and λ, γ, δ in $\mathbb{C}$ (Example 6.2) we have $V(\phi) = \varepsilon\beta$. If

$$U = \frac{\varepsilon\beta}{\varepsilon\beta z_0z_1 + \varepsilon\gamma z_0 + \beta\lambda z_1 + \lambda\gamma}$$

then $V(\phi) = \frac{U}{U \circ \phi}$.

Proposition 6.6. *Let ϕ be an element of $\text{Bir}(\mathbb{P}^3)_{c(\omega)} \setminus \text{Bir}(\mathbb{P}^3)_\omega$. Assume that ϕ preserves a vector field χ non-tangent to ω . Then ϕ preserves a contact form ω' colinear to ω .*

Remark 6.7. Under these assumptions ϕ preserves the vector field χ and the Reeb vector field Z associated to ω' . With the previous notations if $f = z_0\chi_1 + \chi_2$ and $g = z_0Z_1 + Z_2$ one has $V(\phi) = \frac{f \circ \phi}{f} = \frac{g \circ \phi}{g}$. In particular if $H = f/g$ is non-constant, then H is non-constant and invariant: $H \circ \phi = H$.

Proof of Proposition 6.6. Write χ as $\chi_0 \frac{\partial}{\partial z_0} + \chi_1 \frac{\partial}{\partial z_1} + \chi_2 \frac{\partial}{\partial z_2}$ and ϕ as (ϕ_0, ϕ_1, ϕ_2) . Then $\phi_*\chi = \chi$ if and only if $d\phi_i(\chi) = \chi_i \circ \phi$ for $i = 0, 1$ and 2 . Therefore $\phi^*\omega(\chi) = V(\phi)\omega(\chi)$ can be rewritten

$$\phi_0 d\phi_1(\chi) + d\phi_2(\chi) = \phi_0 \chi_1 \circ \phi + \chi_2 \circ \phi = V(\phi)(z_0 \chi_1 + \chi_2).$$

The vector field χ is not tangent to ω , i.e. $\omega(\chi) \neq 0$ or in other words $z_0\chi_1 + \chi_2 \neq 0$ and so

$$V(\phi) = \frac{(z_0\chi_1 + \chi_2) \circ \phi}{z_0\chi_1 + \chi_2}.$$

As a consequence ϕ preserves a contact form ω' colinear to ω (Proposition 6.4). $\square$

Remark 6.8. Let $\phi \in \text{Bir}(\mathbb{P}^3)_{c(\omega)} \setminus \text{Bir}(\mathbb{P}^3)_\omega$. Assume that there exists a vector field χ such that $\phi_*\chi = W\chi$. If W can be written $\frac{U \circ \phi}{U}$, then ϕ preserves the vector field $Y = U\chi$. According to Proposition 6.6 the map ϕ belongs to $\text{Bir}(\mathbb{P}^3)_{\omega'}$ where ω' denotes a contact form colinear to ω .

7. REGULAR BIRATIONAL MAPS

Let $\mathbf{e}_i$ be the point of $\mathbb{P}_{\mathbb{C}}^3$ whose all components are zero except the i -th.

Let us denote by $\mathcal{H}_{\infty}$ the hyperplane $z_3 = 0$. As $\mathcal{H}_{\infty}$ is the unique invariant surface of $c(\omega)$ one has the following statement:

Proposition 7.1. *The hyperplane $\mathcal{H}_{\infty}$ is either preserved, or blown down by any element of $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$.*

Example 7.2. Let ϕ be a birational map of the complex projective plane; $\mathcal{K}(\phi)$ is polynomial if and only if $\phi = (\beta z_1 + \gamma, \delta z_2 + P(z_1))$ with $P \in \mathbb{C}[z_1]$; remark that such a ϕ is a Jonquières polynomial automorphism. In that case

$$\mathcal{K}(\phi) = \left(\frac{1}{\beta} \left(\delta z_0 - \frac{\partial P(z_1)}{\partial z_1} \right), \beta z_1 + \gamma, \delta z_2 + P(z_1) \right).$$

Note that $\deg P = 1$ if and only if $\mathcal{K}(\phi)$ is an automorphism of $\mathbb{P}^3$. If $\deg P > 1$, then $\text{Ind } \mathcal{K}(\phi) = \{z_1 = z_3 = 0\}$ and $\mathcal{H}_{\infty}$ is blown down onto $\mathbf{e}_3$.

Proposition 7.1 naturally implies the following definition. We say that $\phi \in \text{Bir}(\mathbb{P}^3)_{c(\omega)}$ is **regular at infinity** if $\mathcal{H}_{\infty}$ is preserved by ϕ and if $\phi|_{\mathcal{H}_{\infty}}$ is birational. We denote by $\text{Bir}(\mathbb{P}^3)_{c(\omega)}^{\text{reg}}$ (resp. $\text{Bir}(\mathbb{P}^3)_{\omega}^{\text{reg}}$) the set of regular maps at infinity that belong to $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ (resp. $\text{Bir}(\mathbb{P}^3)_{\omega}$).

Example 7.3. Of course the elements of $\text{Aut}(\mathbb{P}^3)_{c(\omega)}$ (Example 6.2) are regular at infinity.

The contact structure is also given in homogeneous coordinates by the 1-form

$$\overline{\omega} = z_0 z_3 dz_1 + z_3^2 dz_2 - (z_0 z_1 + z_2 z_3) dz_3.$$

Let ϕ be an element of $\text{Bir}(\mathbb{P}^3)_{c(\omega)}^{\text{reg}}$; denote by $\overline{\phi}$ its homogeneization. Since $\phi^* \omega = V(\phi) \omega$ one has $\overline{\phi}^* \overline{\omega} = \overline{V(\phi)} \overline{\omega}$ where $\overline{V(\phi)}$ is a homogeneous polynomial. With these notations one can state:

Lemma 7.4. *Let ϕ be a contact birational map of $\mathbb{P}^3$. Assume that ϕ either preserves $\mathcal{H}_{\infty}$, or blows down $\mathcal{H}_{\infty}$ onto a subset contained in $\mathcal{H}_{\infty}$.*

The map ϕ is regular if and only if $\overline{V(\phi)}$ does not vanish identically on $\mathcal{H}_{\infty}$.

Proof. Let us work in the affine chart $z_2 = 1$. On the one hand

$$\overline{\omega} \wedge d\overline{\omega} = -z_3^2 dz_0 \wedge dz_1 \wedge dz_3$$

and on the other hand

$$\phi^*(\overline{\omega} \wedge d\overline{\omega}) = \overline{V(\phi)}^2 \overline{\omega} \wedge d\overline{\omega}.$$

Hence

$$\overline{\phi}_3^2 \det \text{jac } \overline{\phi} = \overline{V(\phi)}^2 z_3^2 \quad (7.1)$$

where $\overline{\phi}_3$ is the third component of $\overline{\phi}$ expressed in the affine chart $z_2 = 1$.

Suppose that ϕ is regular. Let p be a generic point of $\mathcal{H}_{\infty}$. As ϕ is regular, $\overline{\phi}|_{\mathcal{H}_{\infty}}$ is a local diffeomorphism at p . Since $\overline{\phi}$ is birational and p is generic, $\overline{\phi}_p$ is a local diffeomorphism. As a consequence $\det \text{jac } \overline{\phi}$ is an unit at p ; moreover the invariance of $\mathcal{H}_{\infty}$ by $\overline{\phi}$ implies that $\overline{\phi}_3 = z_3 u$ where u is a unit. Therefore $\overline{V(\phi)}$ does not vanish at p .

Conversely assume that $\overline{V(\phi)}$ does not vanish identically on $\mathcal{H}_{\infty}$. As ϕ either preserves $\mathcal{H}_{\infty}$, or contracts $\mathcal{H}_{\infty}$ onto a subset in $\mathcal{H}_{\infty}$, one can write $\overline{\phi}_3$ as $z_3 P$. As a result

$$(7.1) \quad \Leftrightarrow \quad P^2 \det \text{jac } \overline{\phi} = \overline{V(\phi)}^2$$

Since $\overline{V(\phi)}$ does not vanish the map ϕ is then regular at infinity. $\square$

Corollary 7.5. *One has $\text{Bir}(\mathbb{P}^3)_\omega^{\text{reg}} = \text{Aut}(\mathbb{P}^3)_\omega$.*

Proof. Let ϕ be an element of $\text{Bir}(\mathbb{P}^3)_\omega^{\text{reg}}$. From $\phi^*\omega = \omega$, one gets with the previous notations $\bar{\phi}^*\bar{\omega} = z_3^n\bar{\omega}$ for some integer n . Lemma 7.4 implies that $n = 0$, that is $\bar{\phi}^*\bar{\omega} = \bar{\omega}$; then looking at the degree of the members of this equality one gets $\deg \phi = 1$. $\square$

Example 7.6. The group $\text{Bir}(\mathbb{P}^3)_{c(\omega)}^{\text{reg}}$ contains blow-ups in restriction to $\mathcal{H}_\infty$. Indeed let us look at ω in the affine chart $z_2 = 1$ and consider the birational map ϕ given in $z_2 = 1$ by

$$\phi = (z_0, z_0z_1 - z_3, z_0z_3).$$

Since $(\phi^n)^*\omega = z_0^{-n}\omega$, $\phi^n \in \text{Bir}(\mathbb{P}^3)_{c(\omega)}^{\text{reg}} \setminus \text{Bir}(\mathbb{P}^3)_\omega$ for any $n \neq 0$; in restriction to $\mathcal{H}_\infty$ the map ϕ^n coincides with $(z_0, z_1z_0^n)$.

Let us note that $\text{Ind} \phi^n = \{\mathbf{e}_1\} \cup (z_0 = z_2 = 0)$, that $z_0 = 0$ is contracted by ϕ onto $(z_0 = z_2 = 0)$ and $z_2 = 0$ onto $(z_0 = z_3 = 0)$. Besides $\text{Ind} \phi^{-n} = \{z_0 = z_2 = 0\} \cup \{z_0 = z_3 = 0\}$, $(z_0 = 0)$ is blown down by ϕ^{-1} onto $\mathbf{e}_2$ and $(z_2 = 0)$ onto $\mathbf{e}_1$.

Remark 7.7. The group generated by Examples 7.3 and 7.6 is in restriction to $\mathcal{H}_\infty$ and in the affine chart $z_2 = 1$

$$\left\langle \left(\frac{\gamma z_0}{\beta z_1 + \lambda}, \frac{\lambda z_1}{\gamma(\beta z_1 + \lambda)} \right), (z_0, z_0z_1) \mid \gamma, \beta \in \mathbb{C}^*, \lambda \in \mathbb{C} \right\rangle;$$

it is of course a subgroup of $\text{Bir}(\mathbb{P}^3)_{c(\omega)}^{\text{reg}}$.

Question 2. *Does this group coincide with $\text{Bir}(\mathbb{P}^3)_{c(\omega)}^{\text{reg}}$?*

Examples 7.8. a) If ϕ is either a monomial map (i.e. a map of the form $(z_1^p z_2^q, z_1^r z_2^s)$ with $\begin{bmatrix} p & q \\ r & s \end{bmatrix}$ in $\text{GL}(2; \mathbb{Z})$), or a non-linear polynomial automorphism, or a Jonquières map, then $\mathcal{K}(\phi)$ is not regular at infinity.

b) The map of order 5 given by $\left(-\frac{z_2+1+z_0z_1}{z_0z_1^2}, z_2, \frac{z_2+1}{z_1} \right)$, the map $\left(\frac{z_0}{(z_2+1)^2}, z_1, \frac{z_2}{z_2+1} \right)$ and Examples 6.3 a) are non-regular at infinity.

c) Any map of the form

$$\left(\frac{1}{z_0} - f'(z_2), z_2, z_1 + f(z_2) \right)$$

is in $\text{Bir}(\mathbb{P}^3)_{c(\omega)} \setminus \text{Bir}(\mathbb{P}^3)_\omega$ and is not regular at infinity.

d) Elements of the Legendre family are not regular at infinity.

8. EXACT BIRATIONAL MAPS

8.1. First properties. Recall that an element ϕ of $\text{Bir}(\mathbb{P}^2)_\eta$ is **exact** if it can be lifted via ς to $\text{Bir}(\mathbb{P}^3)_\omega$, or equivalently if it belongs to $\text{im } \varsigma$. The following statement allows to determine such maps.

Theorem 8.1. *A map $(\phi_0(z_0, z_1), \phi_1(z_0, z_1)) \in \text{Bir}(\mathbb{P}^2)_\eta$ is exact if and only if the closed form $\phi_0 d\phi_1 - z_0 dz_1$ has trivial residues. In that case $\phi_0 d\phi_1 - z_0 dz_1 = -db$ with $b \in \mathbb{C}(z_0, z_1)$ and*

$$\phi = (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1))$$

belongs to $\text{Bir}(\mathbb{P}^3)_\omega$.

Proof. Remark that $\phi = (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1))$ belongs to $\text{Bir}(\mathbb{P}^3)_{\omega}$ if and only if

$$\phi_0 d\phi_1 - z_0 dz_1 = -db;$$

in other words $\phi_0 d\phi_1 - z_0 dz_1$ is not only a closed rational 1-form but also an exact one. Recall that a closed rational 1-form Θ can be written ([14])

$$\Theta = \sum_i \lambda_i \frac{df_i}{f_i} + dg$$

where the λ_i are complex numbers and the f_i 's and g are rational. The 1-form Θ is exact (i.e. the differential of a rational function) if $\lambda_i = 0$ for all i , that is if the residues of Θ are trivial. $\square$

Example 8.2. The set

$$\left\{ \left(A(z_0), \frac{z_1}{A'(z_0)} \right) \mid A \in \text{PGL}(2; \mathbb{C}) \right\}$$

is a subgroup of exact maps isomorphic to $\text{PGL}(2; \mathbb{C})$; it is a direct consequence of Theorem 8.1.

An other direct consequence of Theorem 8.1 is the following statement:

Corollary 8.3. *The maps $\phi = (\phi_0, \phi_1)$ of $\text{Bir}(\mathbb{P}^2)_{\eta}$ such that $\phi_0 d\phi_1 - z_0 dz_1$ has trivial residues form a group.*

8.2. Involutions. Bertini gives a classification of birational involutions ([6]): a non-trivial birational involution is conjugate to either a Jonquières involution of degree ≥ 2 , or a Bertini involution, or a Geiser involution. More recently Bayle and Beauville precise it ([5]); the map which associates to a birational involution of $\mathbb{P}^2$ its normalized fixed curve establishes a one-to-one correspondence between:

- conjugacy classes of Jonquières involutions of degree d and isomorphism classes of hyperelliptic curves of genus $d - 2$ ($d \geq 3$);
- conjugacy classes of Geiser involutions and isomorphism classes of non-hyperelliptic curves of genus 3;
- conjugacy classes of Bertini involutions and isomorphism classes of non-hyperelliptic curves of genus 4 whose canonical model lies on a singular quadric.

Besides the Jonquières involutions of degree 2 form one conjugacy class.

Proposition 8.4. *Let $I \in \text{Bir}(\mathbb{P}^2)$ be a birational involution. If I is conjugate to either a Geiser involution, or a Bertini involution, or a Jonquières involution of degree ≥ 3 , then I does not belong to $\text{Bir}(\mathbb{P}^2)_{\eta}$.*

Hence the only involutions in $\text{Bir}(\mathbb{P}^2)_{\eta}$ are birationally conjugate to $(-z_0, -z_1)$. Some of them can not be lifted.

Proof. Let us consider such an involution, then the set of fixed points contains a curve Γ of genus > 0 and thus it is not contained in the line at infinity. The jacobian determinant of I at a fixed point of Γ is -1 hence I does not preserve η .

Contrary to the polynomial case (Proposition 3.8) $\text{Bir}(\mathbb{P}^2)_{\eta}$ contains periodic elements that are non-exact. Consider the map $(\phi_0(z_0, z_1), \phi_1(z_0, z_1))$ where

$$\phi_0(z_0, z_1) = -z_0 + \frac{1}{z_1^2 - 1}, \quad \phi_1(z_0, z_1) = -z_1;$$

it is a birational involution that preserves η . Furthermore the 1-form $\phi_0 d\phi_1 - z_0 dz_1$ has non-trivial residues and so is not exact (Theorem 8.1). $\square$

8.3. Quadratic maps. Any birational map of $\mathbb{P}^2$ can be written as a composition of birational maps of degree ≤ 2 (see for instance [1]). The three following maps are birational and of degree 2

$$\begin{aligned}\sigma: \mathbb{P}^2 &\dashrightarrow \mathbb{P}^2 & (z_0 : z_1 : z_2) &\dashrightarrow (z_1 z_2 : z_0 z_2 : z_0 z_1) \\ \rho: \mathbb{P}^2 &\dashrightarrow \mathbb{P}^2 & (z_0 : z_1 : z_2) &\dashrightarrow (z_0 z_2 : z_0 z_1 : z_2^2) \\ \tau: \mathbb{P}^2 &\dashrightarrow \mathbb{P}^2 & (z_0 : z_1 : z_2) &\dashrightarrow (z_0 z_2 + z_1^2 : z_1 z_2 : z_2^2)\end{aligned}$$

Denote by $\mathring{\text{Bir}}_2(\mathbb{P}^2)$ the set of birational maps of $\mathbb{P}^2$ of degree 2 exactly; for any $\phi \in \text{Bir}(\mathbb{P}^2)$ set

$$O(\phi) = \{g\phi h^{-1} \mid g, h \in \text{Aut}(\mathbb{P}^2)\}$$

one has ([13])

$$\mathring{\text{Bir}}_2(\mathbb{P}^2) = O(\sigma) \cup O(\rho) \cup O(\tau).$$

Let us now describe the quadratic birational maps that preserve η ; note that τ preserves η . Consider Y the set of pairs $(g(\gamma), g(\beta))$ where

$$g(\beta) = \left(\frac{\beta_0 z_0 + \beta_1 z_1 + \beta_2}{\beta_6 z_0 + \beta_7 z_1 + \beta_8}, \frac{\beta_3 z_0 + \beta_4 z_1 + \beta_5}{\beta_6 z_0 + \beta_7 z_1 + \beta_8} \right)$$

in $\text{Aut}(\mathbb{P}^2) \times \text{Aut}(\mathbb{P}^2)$ such that

$$\gamma_6 = 0, \quad \gamma_7 \beta_3 = 0, \quad \gamma_7 \beta_4 = 0, \quad \det g \det h = (\gamma_7 \beta_5 + \gamma_8)^3.$$

Proposition 8.5. *A quadratic birational map that preserves η belongs to $O(\tau)$.*

More precisely a birational map belongs to $\mathring{\text{Bir}}_2(\mathbb{P}^2) \cap \text{Bir}(\mathbb{P}^2)_\eta$ if and only if it can be written $g(z_0 + z_1^2, z_1)h$ with (g, h) in Y .

Proof. Let ψ be in $\text{Bir}(\mathbb{P}^2)_\eta \cap \mathring{\text{Bir}}_2(\mathbb{P}^2)$; it is sufficient to prove that $\psi \notin O(\sigma) \cup O(\rho)$.

Assume by contradiction that ψ belongs to $O(\sigma)$, i.e. $\psi = g\sigma h$ with $g = g(\gamma)$, $h^{-1} = g(\beta)$. One can rewrite $\psi^*\eta = \eta$ as $\sigma^*g^*\eta = h^*\eta$; this last one relation is equivalent in the affine chart $z_3 = 1$ to

$$\frac{(\det g) z_0 z_1}{(\gamma_6 z_1 + \gamma_7 z_0 + \gamma_8 z_0 z_1)^3} \eta = \frac{\det h}{(\beta_6 z_0 + \beta_7 z_1 + \beta_8)^3} \eta \quad (8.1)$$

the coefficients γ_6 and γ_7 have thus to be zero and (8.1) is equivalent to

$$\frac{\det g}{\gamma_8^3 z_0^2 z_1^2} \eta = \frac{\det h}{(\beta_6 z_0 + \beta_7 z_1 + \beta_8)^3} \eta$$

and this equality never holds.

A similar argument allows to exclude the case: $\psi \in O(\rho)$. This proves the first assertion.

Let us consider $\psi = g\tau h$ in $\mathring{\text{Bir}}_2(\mathbb{P}^2) \cap \text{Bir}(\mathbb{P}^2)_\eta$ with $g = g(\gamma)$ and $h = g(\beta)$. The 1-form η has a line of poles of order 3 at infinity so does $\psi^*\eta$ and so does $(z_0 + z_1^2, z_1)^*g^*\eta$. But

$$(z_0 + z_1^2, z_1)^*g^*\eta = \frac{\det g}{(\gamma_6(z_0 + z_1^2) + \gamma_7 z_1 + \gamma_8)^3} \eta$$

therefore γ_6 has to be 0. This implies that

$$\psi^*\eta = \frac{\det g \det h}{(\gamma_7(\beta_3 z_0 + \beta_4 z_1 + \beta_5) + \gamma_8)^3} \eta$$

as a consequence $\psi^*\eta = \eta$ if and only if

$$\gamma_6 = 0, \quad \gamma_7\beta_3 = 0, \quad \gamma_7\beta_4 = 0, \quad \det g \det h = (\gamma_7\beta_5 + \gamma_8)^3.$$

□

Theorem 8.6. *A generic element of $\mathring{\text{Bir}}_2(\mathbb{P}^2) \cap \text{Bir}(\mathbb{P}^2)_{\eta}$ is not exact.*

In fact there exists a non-empty Zariski open subset $\tilde{Y}$ of Y such that no element of

$$\{g(\gamma) \tau g(\beta) \mid (g(\gamma), g(\beta)) \in \tilde{Y}\}$$

is exact.

Proof. It is sufficient to exhibit a non-exact element. Let us recall that the birational map $\phi = (\phi_0, \phi_1)$ belongs to $\mathring{\text{Bir}}_2(\mathbb{P}^2) \cap \text{Bir}(\mathbb{P}^2)_{\eta}$ if and only if it can be written as $g(\gamma) \tau g(\beta)$ with $(g(\gamma), g(\beta))$ in Y (Proposition 8.5).

If we consider the special case $\gamma_i = \beta_i = 0$ for any $i \in \{1, 2, 3, 4, 6, 8\}$, $\gamma_5 = \gamma_7$ and $\gamma_0 = \frac{\gamma_7\beta_5^2}{\beta_0\beta_7}$ then

$$z_0 dz_1 - \phi_0 d\phi_1 = -\frac{\beta_5^2 dz_1}{\beta_0 \beta_7 z_1}$$

But $\det g(\beta) \neq 0$ so $\beta_5 \neq 0$ and ϕ can not be lifted to $\text{Bir}(\mathbb{P}^3)_{\omega}$.

The set Y is rational hence irreducible, this yields the result. □

8.4. Examples of exact maps.

Proposition 8.7. *Let ϕ be an automorphism of $\mathbb{P}^2$; the map ϕ is exact if and only if ϕ is affine in the affine chart $z_2 = 1$ and preserves η , that is*

$$\phi = (\delta_0 z_0 + \beta_0 z_1 + \gamma_0, \delta_1 z_0 + \beta_1 z_1 + \gamma_1)$$

with $\delta_i, \beta_i, \gamma_i$ in $\mathbb{C}$ such that $\delta_0\beta_1 - \delta_1\beta_0 = 1$.

Proof. The form η has a pole at infinity so if $\phi \in \text{Aut}(\mathbb{P}^2)$ preserves η , it preserves the pole. Hence ϕ belongs to Aff_2 , so in particular to $\text{Aut}(\mathbb{C}^2)_{\eta}$ and then ϕ is exact. □

We will now consider the subgroup of $\text{Bir}(\mathbb{P}^2)_{\eta}$ that preserves the fibration $z_0 z_1 = \text{cst}$ fiberwise. The following statement says that this subgroup is not isomorphic to the subgroup of $\text{Bir}(\mathbb{P}^2)_{\eta}$ that preserves $z_1 = \text{cst}$ fiberwise.

Proposition 8.8. *The set*

$$\Lambda = \left\{ \left(z_0 a(z_0 z_1), \frac{z_1}{a(z_0 z_1)} \right) \mid a \in \mathbb{C}(t) \right\}$$

is a subgroup isomorphic to the uncountable abelian subgroup $\{(a(z_1)z_0, z_1) \mid a \in \mathbb{C}(z_1)^*\}$ and is contained in $\text{Bir}(\mathbb{P}^2)_{\eta}$.

Any birational map of the form $\left(z_0 a(z_0, z_1), \frac{z_1}{a(z_0, z_1)} \right)$ that preserves η belongs to Λ .

A generic element of Λ is in $\text{Bir}(\mathbb{P}^2)_{\eta}$ but not in $\text{im } \varsigma$. More precisely $\left(z_0 a(z_0 z_1), \frac{z_1}{a(z_0 z_1)} \right) \in \Lambda$ is exact if and only if a is a monomial.

If a is a monomial, i.e. $a(z_0 z_1) = cz_0^{\mu} z_1^{\mu}$ with $c \in \mathbb{C}^*$ and $\mu \in \mathbb{Z}$, then the ς -lifted maps are

$$\left(z_0 cz_0^{\mu} z_1^{\mu}, \frac{z_1}{cz_0^{\mu} z_1^{\mu}}, z_2 - \mu z_0 z_1 + \beta \right), \quad \beta \in \mathbb{C}$$

These maps form a subgroup of $\text{Bir}(\mathbb{P}^3)_\omega$ isomorphic to $\mathbb{C} \times \mathbb{C}^* \times \mathbb{Z}$.

Proof. The first assertion follows from

$$\left(z_0 a(z_0 z_1), \frac{z_1}{a(z_0 z_1)} \right) = (z_0, z_0 z_1)^{-1} (z_0 a(z_1), z_1) (z_0, z_0 z_1)$$

A direct computation shows that $\Lambda \subset \text{Bir}(\mathbb{P}^2)_\eta$.

A birational map $\left(z_0 a(z_0, z_1), \frac{z_1}{a(z_0, z_1)} \right)$ preserves η if and only if

$$\left(z_0 \frac{\partial}{\partial z_0} - z_1 \frac{\partial}{\partial z_1} \right) (a) = 0$$

that is, if and only if $a = a(z_0 z_1)$.

Let us consider $\phi = (\phi_0, \phi_1) = \left(z_0 a(z_0 z_1), \frac{z_1}{a(z_0 z_1)} \right)$ an element of Λ ; then

$$\phi_0 d\phi_1 - z_0 dz_1 = t \frac{a'(t)}{a(t)} dt$$

with $t = z_0 z_1$. Let us write a as follows:

$$a(t) = \prod_{i=1}^n (t - t_i)^{\mu_i}$$

then

$$t \frac{a'(t)}{a(t)} dt = t \sum_{i=1}^n \frac{\mu_i}{t - t_i} dt$$

and the residues of this 1-form are trivial if and only if a is monomial, i.e. $a(t) = ct^\mu$ where $c \in \mathbb{C}^*$ and $\mu \in \mathbb{Z}$. $\square$

We can determine $\mathcal{J} \cap \text{Bir}(\mathbb{P}^2)_\eta$ and the exact maps in $\mathcal{J} \cap \text{Bir}(\mathbb{P}^2)_\eta$.

Proposition 8.9. *A Jonquière map of $\mathbb{P}^2$ preserves η if and only if it can be written as follows*

$$\left(\frac{(\gamma z_1 + \delta)^2}{\varepsilon \delta - \beta \gamma} z_0 + r(z_1), \frac{\varepsilon z_1 + \beta}{\gamma z_1 + \delta} \right)$$

where r belongs to $\mathbb{C}(z_1)$ and $\begin{bmatrix} \varepsilon & \beta \\ \gamma & \delta \end{bmatrix}$ to $\text{PGL}(2; \mathbb{C})$.

Furthermore it is exact if it has the following form

$$\left(\frac{(\gamma z_1 + \delta)^2}{\varepsilon \delta - \beta \gamma} z_0 + P(z_1)(\gamma z_1 + \delta)^2, \frac{\varepsilon z_1 + \beta}{\gamma z_1 + \delta} \right)$$

where P denotes an element of $\mathbb{C}[z_1]$.

Let us now look at monomial maps that belong to $\text{Bir}(\mathbb{P}^2)_\eta$ and those who are exact.

Proposition 8.10. *A monomial map belongs to $\text{Bir}(\mathbb{P}^2)_\eta$ if and only if it can be written either*

$$\left(\gamma z_0^p z_1^{p-1}, \frac{1}{\gamma} z_0^{1-p} z_1^{2-p} \right) \tag{8.2}$$

or

$$\left(\gamma z_0^p z_1^{p+1}, -\frac{1}{\gamma} z_0^{1-p} z_1^{-p} \right) \tag{8.3}$$

with γ in $\mathbb{C}^*$ and p in $\mathbb{Z}$.

Furthermore any monomial map of $\text{Bir}(\mathbb{P}^2)_{\eta}$ is exact.

The ς -lifts of a map of type (8.2) are

$$\left(\gamma z_0^p z_1^{p-1}, \frac{1}{\gamma} z_0^{1-p} z_1^{2-p}, z_2 + (p-1)z_0 z_1 + \beta \right) \quad \beta \in \mathbb{C}$$

similarly the ς -lifts of a map of type (8.3) are

$$\left(\gamma z_0^p z_1^{p+1}, -\frac{1}{\gamma} z_0^{1-p} z_1^{-p}, z_2 + (1-p)z_0 z_1 + \beta' \right) \quad \beta' \in \mathbb{C}$$

Remarks 8.11. – Both maps of type (8.2) and of type (8.3) preserve $(z_0 z_1)^2 = \text{cst}$.

- Maps of type (8.2) form a group G_1 . Note that the matrices $\begin{bmatrix} p & p-1 \\ 1-p & 2-p \end{bmatrix}$ are in $\text{SL}(2; \mathbb{Z})$; they are stochastic up to transposition and have trace equal to 2. The group

$$\left\{ \begin{bmatrix} p & p-1 \\ 1-p & 2-p \end{bmatrix} \mid p \in \mathbb{Z} \right\}$$

is isomorphic to $\mathbb{Z}$. As a consequence G_1 is isomorphic to $\mathbb{C}^* \times \mathbb{Z}$.

The maps of type (8.3) don't form a group. The corresponding matrices $\begin{bmatrix} p & p+1 \\ 1-p & -p \end{bmatrix}$ have determinant -1 , trace 0 and are stochastic up to transposition.

But the union of the maps of type (8.2) or (8.3) is a group which is a double extension of $\mathbb{C}^* \times \mathbb{Z}$.

9. INDETERMINACY AND EXCEPTIONAL SETS

As we have seen if ϕ is a contact map, then $\mathcal{H}_{\infty}$ is either preserved by ϕ , or blown down by ϕ (Proposition 7.1). In case it is blown down, $\mathcal{H}_{\infty}$ can be blown down onto a point or onto a curve; in this last eventuality $\mathcal{H}_{\infty}$ can be contracted onto a curve contained in $\mathcal{H}_{\infty}$ (take for instance $\phi = \mathcal{K}(z_1, z_1 z_2)$). Note also that $\mathcal{H}_{\infty}$ can be contracted onto a curve not contained in $\mathcal{H}_{\infty}$: the map $\mathcal{K}\left(\frac{z_1}{z_2}, \frac{1}{z_2}\right)$ blows down $\mathcal{H}_{\infty}$ onto the legndrian curve $z_0 = z_2 = 0$. We will see that this is a general case and for any contracted surface:

Proposition 9.1. *Let ϕ be a contact birational map of $\mathbb{P}^3$. Assume that ϕ blows down a surface S onto a curve C . Then*

- either C is contained in $\mathcal{H}_{\infty}$,
- or C is an algebraic legndrian curve.

Corollary 9.2. *Let ϕ be a contact birational map of $\mathbb{P}^3$. If C is a curve not contained in $\mathcal{H}_{\infty}$ and blown-up by ϕ on a surface distinct from $\mathcal{H}_{\infty}$, then C is a legndrian curve.*

Let us now give an example of maps of finite order that illustrates Proposition 9.6.

Example 9.3. Start with the birational map $\varphi = \left(z_2, \frac{z_2+1}{z_1} \right)$ of order 5. The map $\mathcal{K}(\varphi) = \left(-\frac{z_2+1+z_0 z_1}{z_0 z_1^2}, z_2, \frac{z_2+1}{z_1} \right)$ blows down $z_2 = -z_3$ onto the legndrian curve $(z_2 = z_1 + z_3 = 0)$;

Proof of Proposition 9.1. We will distinguish the cases $S = \mathcal{H}_{\infty}$ and $S \neq \mathcal{H}_{\infty}$.

Let us start with the eventuality $S = \mathcal{H}_{\infty}$. Suppose that C is not contained in $\mathcal{H}_{\infty}$. Note that $\phi|_{\mathcal{H}_{\infty} \setminus \text{Ind}\phi}$ is holomorphic of rank ≤ 1 . If p belongs to $C \setminus \text{Ind}\phi$, then $\phi^{-1}(p)$ is a curve contained in $\mathcal{H}_{\infty}$; there exists a curve C' transverse to

$$\{\phi^{-1}(p) \mid p \in C \setminus \text{Ind}\phi\}$$

contained in $\mathcal{H}_\infty$ and such that $\phi(C') = C$. Consider a parametrization s of C' ; then $t = \phi \circ s$ is a parametrization of C and

$$t^*\omega = (\phi \circ s)^*\omega = s^*\phi^*\omega = s^*V(\phi)\omega = V(\phi) \circ s \cdot s^*\omega = 0.$$

Assume now that $S \neq \mathcal{H}_\infty$ and $C \not\subset \mathcal{H}_\infty$. Set $C = \phi(S)$. Let us consider a generic point p of S . The germ ϕ_p is holomorphic and $\phi(p) \in C$ does not belong to $\mathcal{H}_\infty$. In particular the 3-form $\phi^*\omega \wedge d\omega$ is thus holomorphic at p ; in fact $V(\phi)_p$ is holomorphic and as we have seen

$$\phi^*\omega \wedge d\omega = V(\phi)^2\omega \wedge d\omega.$$

Since S is blown down by ϕ , the jacobian determinant of ϕ is identically zero on S and then $V(\phi)$ vanishes on S .

Assume that C is not a legendrian curve, then the restriction of ω to C in a neighborhood of $\phi(p)$ defines a 1-form Θ on C without zero (let us recall that p is generic). As the restriction

$$\phi_{p|S,p} : S_p \rightarrow C_{\phi(p)}$$

is locally a submersion, $\phi_{p|S,p}^* \Theta$ is a nonzero 1-form on S_p : contradiction with the fact that $\phi_p^*\omega$ vanishes on S_p . $\square$

There is no statement if $\phi \in \text{Bir}(\mathbb{P}^3)_{c(\omega)}$ blows down $\mathcal{H}_\infty$ onto a point. Indeed

$$\mathcal{K}\left(\frac{z_1}{z_2^2}, \frac{z_1}{z_2^3}\right) = \left(\frac{z_2 + 3z_0z_1}{z_2(z_2 - 2z_0z_1)}, \frac{z_1}{z_2^2}, \frac{z_1}{z_2^3}\right)$$

contracts $\mathcal{H}_\infty$ onto $\mathbf{e}_3 \notin \mathcal{H}_\infty$ but $\mathcal{K}(z_1z_2, z_1z_2^2)$ contracts $\mathcal{H}_\infty$ onto $\mathbf{e}_2 \in \mathcal{H}_\infty$. But we get some result when $\phi \in \text{Bir}(\mathbb{P}^3)_{c(\omega)}$ blows down a surface distinct from $\mathcal{H}_\infty$ onto a point.

Definition. Let ϕ be a contact birational map of $\mathbb{P}^3$. Let $S = (f = 0)$ be an irreducible surface blown down by ϕ , and let p be a smooth point of S such that ϕ and $V(\phi)$ are holomorphic at p . The multiplicity of contraction of ϕ at p is the greatest integer n such that f_p^n divides $V(\phi)$. One can check that n is independent on p . The integer n is the **multiplicity of contraction of ϕ on S** .

Remark 9.4. Let ϕ be a contact birational map of $\mathbb{P}^3$. If ϕ is holomorphic at $p \in \mathbb{P}^3 \setminus \mathcal{H}_\infty$, then $V(\phi)$ is too.

Example 9.5. Let us consider the birational map ϕ defined in the affine chart $z_1 = 1$ by

$$\phi = \left(\frac{z_0z_3^2}{(z_2 + z_3)^2}, \frac{z_2z_3}{(z_2 + z_3)}, z_3 \right);$$

in this chart $\omega = dz_2 - \frac{z_0 + z_2z_3}{z_3^2} dz_3$ and one can check that $V(\phi) = \frac{z_3^2}{z_2 + z_3^2}$. Furthermore $\mathcal{H}_\infty$ is blown down by ϕ onto the point $(0, 0, 0)$; the multiplicity of contraction of ϕ on $\mathcal{H}_\infty$ is thus 2.

Proposition 9.6. Let ϕ be a map of $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ and let S be an irreducible surface distinct from $\mathcal{H}_\infty$ blown down by ϕ onto a point p . If the multiplicity of contraction of ϕ on S is 1, then p belongs to $\mathcal{H}_\infty$.

Remark 9.7. As soon as the multiplicity of contraction of ϕ on S is > 1 , the point p can be in $\mathbb{P}^3 \setminus \mathcal{H}_\infty$. Let us consider the map of $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ given in the affine chart $z_3 = 1$ by

$$\left(\frac{z_2(nz_0z_1 - z_2)}{z_2 + (1-n)z_0z_1}, z_1z_2^{n-1}, z_1z_2^n \right)$$

with $n \in \mathbb{Z}$. The surface $z_2 = 0$ is blown down onto $\mathbf{e}_3 \notin \mathcal{H}_\infty$. One can check that $V(\phi) = \frac{z_1z_2^n}{z_2 + (1-n)z_0z_1}$ so the multiplicity of contraction of ϕ on $z_2 = 0$ is n if $n \geq 2$ and 0 otherwise.

Proof of Proposition 9.6. Assume by contradiction that $p = (p_0, p_1, p_2)$ does not belong to $\mathcal{H}_{\infty}$. Let $(f = 0)$ be an equation of S ; as the multiplicity of contraction of ϕ on S is 1 one has $V(\phi) = fV_1$ with $V_1|_S$ generically regular. There exists a point $m \in S$ such that $f_{,m}$ is a submersion and ϕ is holomorphic at m . One has $\phi_{,m} = (p_0 + fA, p_1 + fB, p_2 + fC)$ with A, B, C holomorphic and $\phi_{,m}^* \omega = V(\phi)\omega$ can be rewritten

$$(fA + p_0)(f dB + B df) + (f dC + C df) = fV_1(z_0 dz_1 + dz_2) \quad (9.1)$$

This implies that there exists C_1 holomorphic such that $p_0 B + C = fC_1$, i.e. $C = fC_1 - p_0 B$. Hence

$$(9.1) \iff fAdB + ABdf + fdC_1 + 2C_1 df = V_1(z_0 dz_1 + dz_2) \quad (9.2)$$

The multiplicity of contraction of ϕ on S is 1 hence f does not divide V_1 . Then S is invariant by ω and this gives a contradiction with the fact that $\mathcal{H}_{\infty}$ is the only invariant surface of ω . $\square$

For elements in $\text{Bir}(\mathbb{P}^3)_{\omega}$ we only have one statement that includes both cases of a surface contracted onto a point and onto a curve. Let us remark that in the case of a point, we don't need the assumption about the multiplicity of contraction; in the other one the statement shows that Proposition 9.1 applies to elements of $\text{Bir}(\mathbb{P}^3)_{c(\omega)} \setminus \text{Bir}(\mathbb{P}^3)_{\omega}$.

Proposition 9.8. *Let ϕ be a map of $\text{Bir}(\mathbb{P}^3)_{\omega}$. If S is a surface distinct from $\mathcal{H}_{\infty}$ contracted by ϕ , then $\phi(S)$ belongs to $\mathcal{H}_{\infty}$.*

Proof. From $\phi^* \omega = \omega$ one gets $\phi^*(\omega \wedge d\omega) = \omega \wedge d\omega = dz_0 \wedge dz_1 \wedge dz_2$. Suppose that for $p \in S$ generic $\phi(p)$ does not belong to $\mathcal{H}_{\infty}$. As $\text{codim Ind } \phi \geq 2$, the map ϕ is holomorphic at p . Since ϕ preserves the volume form, ϕ is a diffeomorphism; hence ϕ cannot blow down a subvariety onto a curve or a point not contained in $\mathcal{H}_{\infty}$. $\square$

Example 9.9. If $\phi = (\phi_1, \phi_2) = (z_1^p z_2^q, z_1^r z_2^s)$, with $\begin{bmatrix} p & q \\ r & s \end{bmatrix} \in \text{SL}(2; \mathbb{Z})$, then

$$\mathcal{K}(\phi) = \left(z_1^{r-p} z_2^{s-q} \frac{-rz_2 + sz_0 z_1}{pz_2 - qz_0 z_1}, z_1^p z_2^q, z_1^r z_2^s \right).$$

Note that for any $\begin{bmatrix} p & q \\ r & s \end{bmatrix} \in \text{SL}(2; \mathbb{Z})$ the map $\mathcal{K}(\phi)$ belongs to $\text{Bir}(\mathbb{P}^3)_{c(\omega)} \setminus \text{Bir}(\mathbb{P}^3)_{\omega}$.

For instance if $\begin{bmatrix} p & q \\ r & s \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$, i.e. if $\sigma = \left(\frac{1}{z_0}, \frac{1}{z_1} \right)$ is the Cremona involution, then

$$\mathcal{K}(\sigma) = \mathcal{K}(\sigma^{-1}) = \left(\frac{z_0 z_1^2}{z_2^2}, \frac{1}{z_1}, \frac{1}{z_2} \right)$$

and $\text{Ind } \mathcal{K}(\sigma) = \{z_0 = z_2 = 0\} \cup \{z_0 = z_3 = 0\} \cup \{z_1 = z_2 = 0\} \cup \{z_1 = z_3 = 0\}$; furthermore $z_2 = 0$ and $\mathcal{H}_{\infty}$ are blown down onto $\mathbf{e}_1$ and $z_1 = 0$ onto $\mathbf{e}_2$.

Part 3. Some common properties

10. INVARIANT CURVES AND SURFACES

The following statement is a local statement of contact analytic geometry.

Proposition 10.1. *Let ϕ be an element of $\text{Aut}(\mathbb{C}^3)_{\omega}$ or $\text{Bir}(\mathbb{P}^3)_{\omega}$. Suppose that m is a periodic point of ϕ and that there exists a germ of irreducible curve C invariant by ϕ , passing through m . Then*

- either C is a curve of periodic points (i.e. $\phi|_C^{\ell} = \text{id}$ for some integer ℓ),
- or C is a legendrian curve.

Let us note that according to Proposition 11.4 we know that such a situation often occurs.

Proof. Assume that ϕ belongs to $\text{Aut}(\mathbb{C}^3)_\omega$. Up to considering a well-chosen iterate of ϕ let us assume that m is a fixed point of ϕ . Let $s \mapsto \gamma(s)$ be a local parametrization of C at m . Up to reparametrization one can suppose that $\gamma(0) = m$. Let φ be the "restriction" to C of ϕ , that is the local map $\varphi: \mathbb{C}_{,0} \rightarrow \mathbb{C}$ defined by $\varphi(0) = 0$ and

$$\forall s \in \mathbb{C}_{,0} \quad \phi(\gamma(s)) = \gamma(\varphi(s)).$$

On the one hand $\gamma^* \omega = \varepsilon(s)ds$ and on the other hand $\gamma^* \omega = \gamma^* \phi^* \omega = (\phi \circ \gamma)^* \omega$ so

$$\varepsilon(s)ds = \varphi^*(\varepsilon(s)ds) = \varepsilon(\varphi) \varphi' ds.$$

Let us set $\tilde{\varepsilon}(s) = \int_0^s \varepsilon(t)dt$. One has $(\tilde{\varepsilon}(\varphi))' = \varepsilon(\varphi) \varphi' = \varepsilon(s) = (\tilde{\varepsilon}(s))'$ hence $\tilde{\varepsilon}(\varphi) = \tilde{\varepsilon} + \beta$ for some $\beta \in \mathbb{C}$. As $\varphi(0) = 0$, one gets $\beta = 0$ and $\tilde{\varepsilon}(\varphi) = \tilde{\varepsilon}$. Then:

- either $\tilde{\varepsilon} = 0$ therefore $\varepsilon = 0$ and C is a legendrian curve.
- or there exists some local coordinate for which $\tilde{\varepsilon} = z^\ell$, $\varphi = e^{2\pi i k/\ell} z$ and $\phi|_C = \text{id}$.

□

If φ is a polynomial automorphism of $\mathbb{C}^2$ that preserves a curve distinct from the line at infinity, then φ is conjugate to a Jonquière's polynomial automorphism ([9]); in particular φ preserves a rational fibration. We have a similar statement in dimension 3:

Proposition 10.2. *If $\phi \in \text{Aut}(\mathbb{C}^3)_\omega$ preserves a surface, then*

$$\phi = (\varphi(z_0, z_1), z_2 + b(z_0, z_1))$$

where φ is $\text{Aut}(\mathbb{C}^2)$ -conjugate to a Jonquière's polynomial automorphism.

Proof. Let us write ϕ as $(\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1))$ and set $\varphi = (\phi_0, \phi_1)$.

First note that if $b \equiv 0$ then $\phi_0 d\phi_1 - z_0 dz_1 = 0$; as a result $\phi_1 = \phi_1(z_1)$ and φ is a Jonquière's polynomial automorphism.

Let us now assume that the surface S preserved by ϕ is described by

$$a_\ell(z_0, z_1)z_2^\ell + a_{\ell-1}(z_0, z_1)z_2^{\ell-1} + a_{\ell-2}(z_0, z_1)z_2^{\ell-2} + \dots = 0$$

where $a_i \in \mathbb{C}[z_0, z_1]$, or equivalently by

$$z_2^\ell + \tilde{a}_{\ell-1}(z_0, z_1)z_2^{\ell-1} + \tilde{a}_{\ell-2}(z_0, z_1)z_2^{\ell-2} + \dots = 0$$

where $\tilde{a}_i = a_i/a_\ell$. Writing that S is invariant by ϕ one gets that

$$\begin{aligned} & (z_2 + b(z_0, z_1))^\ell + \tilde{a}_{\ell-1}(\varphi(z_0, z_1))(z_2 + b(z_0, z_1))^{\ell-1} + \tilde{a}_{\ell-2}(\varphi(z_0, z_1))(z_2 + b(z_0, z_1))^{\ell-2} + \dots \\ &= z_2^\ell + \tilde{a}_{\ell-1}(z_0, z_1)z_2^{\ell-1} + \tilde{a}_{\ell-2}(z_0, z_1)z_2^{\ell-2} + \dots \end{aligned}$$

Looking at terms in $z_2^{\ell-1}$ one gets that $\ell b(z_0, z_1) = \tilde{a}_{\ell-1}(z_0, z_1) - \tilde{a}_{\ell-1}(\varphi(z_0, z_1))$.

- If $\tilde{a}_{\ell-1}$ is constant, then $b \equiv 0$ and as we just see φ is a Jonquière's polynomial automorphism.
- Otherwise ϕ is conjugate (in $\text{Bir}(\mathbb{P}^3)$) via $\left(z_0, z_1, z_2 + \frac{\tilde{a}_{\ell-1}}{\ell}\right)$ to $\psi = (\varphi, z_2)$. The map ψ preserves $\tilde{\omega} = z_0 dz_1 + d\left(z_2 + \frac{\tilde{a}_{\ell-1}}{\ell}\right)$, the surface $\tilde{S}$ given by

$$z_2^\ell + \tilde{a}_{\ell-2}(z_0, z_1)z_2^{\ell-2} + \tilde{a}_{\ell-3}(z_0, z_1)z_2^{\ell-3} + \dots = 0$$

and thus $\tilde{a}_i(\varphi) = \tilde{a}_i$. If one of the $\tilde{a}_i$ is non-constant, then φ is a Jonquière's polynomial automorphism. Otherwise $\tilde{S} = \cup_j (z_2 = c_j)$; up to take an iterate ψ^k of ψ one can suppose that any $z_2 = c_j$ is invariant. Consider $z_2 = c_0$; up to a well-chosen translation (that belongs to $\text{Bir}(\mathbb{P}^3)_\omega$) the hypersurface $z_2 = 0$ is invariant, that is ψ^k is a Jonquière's map and so does ψ . □

Example 10.3. For any $n \geq 1$ consider $\phi = \left(z_0 + z_1^n, z_1, z_2 - \frac{z_1^{n+1}}{n+1} \right)$ in $\text{Aut}(\mathbb{C}^3)_{\omega}$. The map $\varphi = (z_0 + z_1^n, z_1)$ is a Jonquières polynomial automorphism. The surface S given by $z_2 + \frac{z_0 z_1}{n+1} = 0$, is invariant by ϕ . The foliation induced by ω on S is described by the linear differential equation $nz_0 dz_1 - z_1 dz_0$. In fact the functions $z_2 + \frac{z_0 z_1}{n+1}$ and z_1 are invariant by ϕ and the commutative Lie algebra generated by the vector fields $\frac{\partial}{\partial z_0} + \frac{z_1}{n+1} \frac{\partial}{\partial z_2}$ and $\frac{\partial}{\partial z_2}$ are invariant by ϕ .

In general an element of $\text{Aut}(\mathbb{C}^3)_{\omega}$ has no invariant surface. For instance there is no polynomial solution to

$$-a(\varphi(z_0, z_1)) + a(z_0, z_1) = -\frac{z_1^{n+1}}{n+1} + \beta$$

with $\varphi = (z_0 + z_1^n, z_1)$ as soon as $\beta \neq 0$.

Remark 10.4. If $\phi \in \text{Bir}(\mathbb{P}^3)_{\omega}$ preserves $z_2 = 0$, then ϕ belongs to the Klein family; more precisely $\phi = \left(\frac{z_0}{v(z_1)}, v(z_1), z_2 \right)$ with $v \in \text{PGL}(2; \mathbb{C}(z_1))$. Indeed since ϕ belongs to $\text{Bir}(\mathbb{P}^3)_{\omega}$,

$$\phi = (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1)).$$

But ϕ preserves $z_2 = 0$ so $b \equiv 0$ and $\phi^* \omega = \omega$ implies that $\phi_1 = v(z_1)$ with $v \in \text{PGL}(2; \mathbb{C}(z_1))$ and $\phi_0 = \frac{z_0}{v(z_1)}$.

Of course there are more general contact maps that preserve $z_2 = 0$; let us give some examples:

$$\mathcal{K} \left(z_1, \frac{z_2}{a(z_1)z_2 + 1} \right), \quad \mathcal{K}(z_1 + P(z_2), z_2)$$

where $a \in \mathbb{C}(z_1)^*$ and $P \in \mathbb{C}[z_1]$.

Let ϕ be an element of $\text{Bir}(\mathbb{P}^3)_{\omega}$. Suppose that ϕ preserves a surface S distinct from $\mathcal{H}_{\infty}$. The contact form is non-zero on S so induces a foliation $\mathcal{F}$ on S , necessarily invariant by ϕ ; let us describe $(S, \phi|_S, \mathcal{F})$:

Proposition 10.5. *Let ϕ be an element of $\text{Bir}(\mathbb{P}^3)_{\omega}$ that preserves a surface distinct from $\mathcal{H}_{\infty}$. Then ϕ is $\text{Bir}(\mathbb{P}^3)$ -conjugate to $(\varphi(z_0, z_1), z_2)$ with φ in $\text{Bir}(\mathbb{P}^2)$. The map φ preserves a codimension 1 foliation given by a closed 1-form. As a consequence ϕ preserves a "vertical" foliation and a rational function $z_2 + a(z_0, z_1)$.*

Proof. Let us denote by S the surface invariant by $\phi = (\varphi(z_0, z_1), z_2 + b(z_0, z_1))$ with $\varphi \in \text{Bir}(\mathbb{P}^2)$. One can assume that S is given by

$$z_2^{\ell} + a_{\ell-1}(z_0, z_1)z_2^{\ell-1} + \dots = 0$$

The fact that S is invariant by ϕ implies that $a_{\ell-1}(z_0, z_1) - a_{\ell-1}(\varphi(z_0, z_1)) = \ell b(z_0, z_1)$. Let us consider the map $\psi = \left(z_0, z_1, z_2 + \frac{a_{\ell-1}(z_0, z_1)}{\ell} \right)$. One has

$$\tilde{\phi} = \psi \phi \psi^{-1} = \left(\varphi(z_0, z_1), z_2 + b(z_0, z_1) - \frac{a_{\ell-1}(z_0, z_1)}{\ell} + \frac{a_{\ell-1}(\varphi(z_0, z_1))}{\ell} \right) = (\varphi(z_0, z_1), z_2)$$

As S and ω are invariant by ϕ , the restriction $\phi|_S$ preserves the foliation induced by ω on S , and $\tilde{\phi}$ preserves the "vertical" foliation given by $z_0 dz_1 - da_{\ell-1}(z_0, z_1)$. Therefore φ preserves a codimension 1 foliation given by a closed 1-form. $\square$

Example 10.6. If $\phi = (z_2, z_1 z_2^n)$, then $\mathcal{K}(\phi) = \left(-\frac{z_2^n}{z_0} + nz_1, z_1 z_2^n, z_2 \right)$ belongs to $\text{Bir}(\mathbb{P}^3)_{\mathcal{C}(\omega)} \setminus \text{Bir}(\mathbb{P}^3)_{\omega}$ preserves the surface $z_1 = 0$ and also $z_2 = \text{cst}$.

11. DYNAMICAL PROPERTIES

11.1. Periodic points. Let ϕ be a birational map of $\mathbb{P}^n$; a point p is a **periodic point** of ϕ of period ℓ if ϕ is holomorphic on a neighborhood of any point of $\{\phi^j(q) \mid j = 0, \dots, \ell - 1\}$ and if $\phi^\ell(q) = q$ and $\phi^j(q) \neq q$ for $1 \leq j \leq \ell - 1$.

Recall that a polynomial automorphism of $\mathbb{C}^2$ of Hénon type (see [20]) has an infinite number of hyperbolic periodic points. For any of these points p of period ℓ_p there exists a stable manifold $W^s(p)$ defined as the set of points that move towards the orbit of p by positive iteration of ϕ^{ℓ_p} ; such a $W^s(p)$ is an immersion from $\mathbb{C}$ to $\mathbb{C}^2$. Remark that even if $W^s(m) \neq W^s(p)$ are different as soon as p and m have distinct orbits one has $\overline{W^s(m)} = \overline{W^s(p)}$. The Julia set of ϕ is the topological boundary of the set of points with bounded positive orbits. One can prove that the Julia set of ϕ is equal to the closure of any of the stable manifold. Hence its topology is very complicated: this set contains an infinite number of immersions of $\mathbb{C}$ and pairwise distinct ([20]).

Example 11.1. Let us consider a polynomial automorphism ϕ of Hénon type given by $\phi = (\beta z_1 + z_0^2, -\gamma z_0)$. A ζ -lift of ϕ to $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ is

$$\phi = \left(\beta z_1 + z_0^2, -\gamma z_0, \gamma \beta z_2 + \gamma \beta z_0 z_1 + \frac{\gamma}{3} z_0^3 \right)$$

Take a periodic point (p_0, p_1) of ϕ of period k ; then as $\phi^k = (\phi^k(z_0, z_1), (\gamma\beta)^k z_2 + f(z_0, z_1))$ one gets, as soon as $\gamma\beta$ is not a root of unity, that there exists p_2 such that $\phi^k(p_0, p_1, p_2) = (p_0, p_1, p_2)$.

More generally, one can state:

Proposition 11.2. Let ϕ the element of $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$ of the following type

$$\phi = (\phi, \det \text{jac} \phi z_2 + b(z_0, z_1))$$

with ϕ in $\text{Bir}(\mathbb{P}^2)$ and b in $\mathbb{C}(z_0, z_1)$.

If $\det \text{jac} \phi$ is not a root of unity, then any periodic point of ϕ can be lifted into a periodic point of ϕ .

Corollary 11.3. Let ϕ be a polynomial automorphism of $\mathbb{C}^2$ of Hénon type. A ζ -lift of ϕ has an infinite number of periodic points that lift the hyperbolic periodic points of ϕ .

Question 3. Let ϕ be a Hénon automorphism and let ϕ be a ζ -lift of ϕ . The closure of the hyperbolic periodic points of ϕ is the Julia set of ϕ ; in particular it is a Cantor set. Is the closure of the set of periodic points of ϕ a Cantor set?

Let us consider a Hénon automorphism $\phi = (\phi_1, \phi_2)$ and let m be an hyperbolic periodic point of ϕ ; then the matrix

$$\begin{bmatrix} -\frac{\partial \phi_2}{\partial z_1} & \frac{\partial \phi_2}{\partial z_2} \\ \frac{\partial \phi_1}{\partial z_1} & -\frac{\partial \phi_1}{\partial z_2} \end{bmatrix}$$

is a non-parabolic one and so $z_0 \mapsto \frac{-\frac{\partial \phi_2}{\partial z_1} + \frac{\partial \phi_2}{\partial z_2} z_0}{\frac{\partial \phi_1}{\partial z_1} - \frac{\partial \phi_1}{\partial z_2} z_0}$ has two fixed points. We can thus state the following:

Proposition 11.4. Let ϕ be an automorphism of $\mathbb{C}^2$ of Hénon type; to any periodic point of period ℓ of ϕ corresponds two periodic points of period ℓ of $\mathcal{K}(\phi) \in \text{Bir}(\mathbb{P}^3)_{c(\omega)}$.

A similar question as Question 3 is the following:

Question 4. Let ϕ be a polynomial automorphism of $\mathbb{C}^2$ of Hénon type; what is the topology of the distribution of periodic points of $\mathcal{K}(\phi)$? Is it a discrete set? Is its closure a Cantor set?

Remark 11.5. Let us consider an element $(\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1))$ of $\text{Bir}(\mathbb{P}^3)_{\omega}$. Then $\phi_t = (\phi_0(z_0, z_1), \phi_1(z_0, z_1), z_2 + b(z_0, z_1) + t)$ belongs to $\text{Bir}(\mathbb{P}^3)_{\omega}$. If $p = (p_0, p_1, p_2)$ is a fixed point of ϕ_t , then (p_0, p_1) is a fixed point of $\phi = (\phi_0, \phi_1)$ and $b(p_0, p_1) + t = 0$. In particular if ϕ only has isolated fixed points (that is ϕ has no curve of fixed points, which is the case in general), then ϕ_t has no fixed points for t generic. Similarly, if ϕ has a countable number of periodic points, then for t generic ϕ_t has no periodic points.

11.2. Degree and degree growths. In the 2-dimensional case, that is if ϕ belongs to $\text{Aut}(\mathbb{C}^2)$, or $\text{Bir}(\mathbb{P}^2)$, then $\deg \phi = \deg \phi^{-1}$. This equality is not true in higher dimension; for instance if

$$\phi = (z_0^2 + z_2^2 + z_1, z_2^2 + z_0, z_2),$$

then $\phi^{-1} = (z_1 - z_2^2, z_0 - (z_1 - z_2^2)^2 - z_2^2, z_2)$. What happens in our context? The equality $\deg \phi = \deg \phi^{-1}$ still does not hold; indeed if $(\phi_0, \phi_1, z_2 + b(z_0, z_1))$ belongs to $\text{Aut}(\mathbb{C}^3)_{\omega}$, then $-db = \phi_0 d\phi_1 - z_0 dz_1$ and $\deg b = \deg \phi_0 + \deg \phi_1$. For instance if $\phi = (z_0 + (z_1^3 - z_0)^2, z_1^3 - z_0, z_2)$, then

$$\phi^{-1} = ((z_0 - z_1^3)^3 - z_1, z_0 - z_1^3, z_2).$$

Hence the degree of the ζ -lifts of ϕ (resp. ϕ^{-1}) is 9 (resp. 8).

Let ϕ and ψ be two birational self-maps of $\mathbb{P}^3$. We will say that **the degree growths of ϕ and ψ are of the same order** if one of the following holds

- $(\deg \phi^n)_n$ and $(\deg \psi^n)_n$ are bounded,
- there exist an integer k such that $\lim_{n \rightarrow +\infty} \frac{\deg \phi^n}{n^k}$ and $\lim_{n \rightarrow +\infty} \frac{\deg \psi^n}{n^k}$ are finite and nonzero,
- $(\deg \phi^n)_n$ and $(\deg \psi^n)_n$ grow exponentially.

Let ϕ be a polynomial automorphism of $\mathbb{C}^2$; let us recall that ϕ has either a bounded growth or an exponential one ([20]). Denote by ϕ a ζ -lift of ϕ to $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$

$$\phi = (\phi, \det \text{jac } \phi z_2 + b(z_0, z_1))$$

Note that b belongs to $\mathbb{C}[z_0, z_1]$ and so $\deg b(\phi^j(z_0, z_1)) \leq \deg b \deg \phi^j$ for any j . Hence

$$\deg \phi^n \leq \deg \phi^n \leq \max(\deg \phi^n, \deg b \deg \phi^{n-1})$$

and

- if $(\deg \phi^n)_n$ is bounded, then $(\deg \phi^n)_n$ is bounded,
- if $(\deg \phi^n)_n$ grows exponentially, then $(\deg \phi^n)_n$ grows exponentially.

Remark that if ψ is a polynomial automorphism of $\mathbb{C}^3$ linear growth is also possible ([8]) and this eventuality does not appear when we look at elements of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$.

In the case of the ζ -lift of an exact element of $\text{Bir}(\mathbb{P}^2)_{\eta}$ we cannot give formula because we are not dealing with polynomials. But the degree growth of a ζ -lift ϕ of an exact element ϕ of $\text{Bir}(\mathbb{P}^2)_{\eta}$ and the degree growth of ϕ are the same. Indeed set $\phi^n = (\phi_{0,n}, \phi_{1,n})$ for any $n \geq 1$. On the one hand

$$\phi^n = (\phi_{0,n}, \phi_{1,n}, z_2 + b(z_0, z_1) + b(\phi_{0,1}, \phi_{1,1}) + b(\phi_{0,2}, \phi_{1,2}) + \dots + b(\phi_{0,n-1}, \phi_{1,n-1}))$$

with $db = z_0 dz_1 - \phi_0 d\phi_1$, but on the other hand $\phi^n = (\phi_{0,n}, \phi_{1,n}, z_2 + \tilde{b}(z_0, z_1))$ with $d\tilde{b} = z_0 dz_1 - \phi_{0,n} d\phi_{1,n}$. Using this last writing one gets the statement.

Let ϕ be a birational self-map of $\mathbb{P}^2$. For any $n \geq 1$ set $\phi^n = (\phi_{1,n}, \phi_{2,n}) = \left(\frac{P_{1,n}}{Q_{1,n}}, \frac{P_{2,n}}{Q_{2,n}}\right)$ with $P_{i,n}, Q_{i,n} \in \mathbb{C}[z_0, z_1]$ without common factor; denote by $p_{i,q}$ (resp. $q_{i,n}$) the degree of $P_{i,n}$ (resp. $Q_{i,n}$). Of course

$\deg \phi^n = \max(p_{1,n} + q_{2,n}, p_{2,n} + q_{1,n}, q_{1,n} + q_{2,n})$ and since

$$\mathcal{K}(\phi)^n = \mathcal{K}(\phi^n) = \left(\frac{Q_{2,n}^2}{Q_{1,n}^2} \frac{P_{2,n} \frac{\partial Q_{2,n}}{\partial z_1} - Q_{2,n} \frac{\partial P_{2,n}}{\partial z_1} + \left(Q_{2,n} \frac{\partial P_{2,n}}{\partial z_2} - P_{2,n} \frac{\partial Q_{2,n}}{\partial z_2} \right) z_0}{Q_{1,n} \frac{\partial P_{1,n}}{\partial z_1} - P_{1,n} \frac{\partial Q_{1,n}}{\partial z_1} - \left(Q_{1,n} \frac{\partial P_{1,n}}{\partial z_2} - P_{1,n} \frac{\partial Q_{1,n}}{\partial z_2} \right) z_0}, \frac{P_{1,n}}{Q_{1,n}}, \frac{P_{2,n}}{Q_{2,n}} \right)$$

one gets $\deg \phi^n \leq \deg \mathcal{K}(\phi)^n \leq \max(4q_{2,n} + p_{2,n} + 1, 2p_{1,n} + 2q_{1,n} + q_{2,n} + 1, p_{2,n} + 3q_{1,n} + p_{1,n} + 1)$.

Proposition 11.6. – Assume that $G = \text{Aut}(\mathbb{C}^2)$ or $G = \text{Bir}(\mathbb{P}^2)_\eta$. Let ϕ be an element of G , and let ϕ be a ς -lift of ϕ . The degree growths of ϕ and ϕ are of the same order.

– Let ϕ be a birational self-map of the complex projective plane, and let us consider $\mathcal{K}(\phi)$ the image of ϕ by $\mathcal{K}$. The degree growths of ϕ and $\mathcal{K}(\phi)$ are of the same order.

11.3. Centralisers. If G is a group and f an element of G , we denote by $\text{Cent}(f, G)$ the centraliser of f in G , that is

$$\text{Cent}(f, G) = \{g \in G \mid fg = gf\}$$

Let ϕ be a polynomial automorphism of $\mathbb{C}^2$, then ([20, 29])

- either ϕ is conjugate to an element of J_2 and $\text{Cent}(\phi, \text{Aut}(\mathbb{C}^2))$ is uncountable;
- or ϕ is of Hénon type and the centraliser of ϕ is isomorphic to $\mathbb{Z} \rtimes \mathbb{Z}/p\mathbb{Z}$ for some p .

Let $\mathcal{H}$ be the set of polynomial automorphisms of $\mathbb{C}^2$ of Hénon type.

Proposition 11.7. Let ϕ be a polynomial automorphism of $\mathbb{C}^2$ and let ϕ be one of its ς -lift.

- If $\det \text{jac } \phi = 1$, then $\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_\omega)$ is uncountable and isomorphic to $\text{Cent}(\phi) \rtimes \mathbb{C}$.
- If $\det \text{jac } \phi \neq 1$ and ϕ belongs to $\mathcal{H}$, then $\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})$ is countable and isomorphic to $\text{Cent}(\phi)$.

Proof. One can look at the restriction of ς to $\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})$:

$$\varsigma|_{\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})} : \text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)}) \rightarrow \text{Cent}(\phi, \text{Aut}(\mathbb{C}^2))$$

Of course

$$\ker \varsigma|_{\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})} \subset \{(z_0, z_1, z_2 + \beta) \mid \beta \in \mathbb{C}\}.$$

If $\det \text{jac } \phi = 1$, i.e. ϕ belongs to $\text{Aut}(\mathbb{C}^2)_\eta$, then

$$\ker \varsigma|_{\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})} = \{(z_0, z_1, z_2 + \beta) \mid \beta \in \mathbb{C}\}$$

and the centraliser of a ς -lift of ϕ is always uncountable even if $\text{Cent}(\phi, \text{Aut}(\mathbb{C}^2))$ is countable.

If $\det \text{jac } \phi \neq 1$, i.e. ϕ belongs to $\text{Aut}(\mathbb{C}^2) \setminus \text{Aut}(\mathbb{C}^2)_\eta$, then $\ker \varsigma|_{\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})} = \{\text{id}\}$ and

$$\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)}) \hookrightarrow \text{Cent}(\phi, \text{Aut}(\mathbb{C}^2))$$

In particular if ϕ belongs to $(\text{Aut}(\mathbb{C}^2) \setminus \text{Aut}(\mathbb{C}^2)_\eta) \cap \mathcal{H}$, then $\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})$ is countable. $\square$

Remark 11.8. Contrary to the 2-dimensional case there exist some ϕ in $\text{Aut}(\mathbb{C}^3)_\omega$ such that

- $\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_\omega)$ is uncountable,
- and $(\deg \phi^n)_{n \in \mathbb{N}}$ grows exponentially.

A similar reasoning leads to:

Proposition 11.9. Let $\phi \in \text{Bir}(\mathbb{P}^2)_\eta$ be an exact map, and let ϕ be one of its ς -lifts. Then $\text{Cent}(\phi, \text{Bir}(\mathbb{P}^3)_\omega)$ is uncountable.

Let $G = \text{Aut}(\mathbb{C}^2)$ or $G = \text{Bir}(\mathbb{P}^2)_{\eta}$. Let φ be an element of G , and let ϕ be one of its ς -lift. In the following examples we look at the links between the ς -lift of $\text{Cent}(\varphi, G)$ and $\text{Cent}(\phi, G')$ where $G' = \text{Aut}(\mathbb{C}^3)_{c(\omega)}$ or $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$.

Example 11.10. In this example we give a polynomial automorphism φ and maps in $\text{Cent}(\varphi, \text{Aut}(\mathbb{C}^2))$ whose only one ς -lift belongs to $\text{Aut}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})$ where ϕ denotes a ς -lift of φ .

Let us now consider the Hénon automorphism φ given by

$$\varphi = (\delta z_1, \beta z_1^k - \gamma z_0)$$

where δ, β, γ are complex numbers such that $\delta\beta \neq 0$, $\delta\beta \neq 1$ and $k \geq 4$. The map

$$\phi = \left(\delta z_1, \beta z_1^k - \gamma z_0, \delta \gamma z_2 + \delta \gamma z_0 z_1 - \frac{\delta \beta}{k+1} z_1^{k+1} \right)$$

is a ς -lift of φ . One can check that $(\zeta z_0, \zeta z_1)$, where $\zeta \in \mathbb{C}^*$ such that $\zeta^k = \zeta$, commutes with φ . Among the ς -lifts $(\zeta z_0, \zeta z_1, \zeta^2 z_2 + \beta)$, $\beta \in \mathbb{C}$, only one commutes with ϕ .

Example 11.11. We consider a polynomial automorphism φ , a subgroup G of $\text{Cent}(\varphi, \text{Aut}(\mathbb{C}^2))$ and G_{ς} its ς -lift. In the first example the inclusion $G_{\varsigma} \subset \text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})$ holds whereas in the second example it doesn't.

Let us consider the polynomial automorphism $\varphi = (\beta^d z_0 + \beta^d z_1^d Q(z_1^r), \beta z_1)$ with $\beta \in \mathbb{C}^*$, $Q \in \mathbb{C}[z_1]$ and $d, r \in \mathbb{N}$. One can check that

$$G = \{ (z_0 + \gamma z_1^d, z_1) \mid \gamma \in \mathbb{C} \} \subset \text{Cent}(\varphi, \text{Aut}(\mathbb{C}^2))$$

The map $\phi = (\beta^d z_0 + \beta^d z_1^d Q(z_1^r), \beta z_1, \beta^{d+1} z_2 - \beta P(z_1))$ with $P'(z_1) = z_1^q Q(z_1^r)$ is a ς -lift of φ . Let G_{ς} be the ς -lift of G ; the group

$$G_{\varsigma} = \left\{ \left(z_0 + \gamma z_1^d, z_1, z_2 - \frac{\gamma z_1^{d+1}}{d+1} \right) \mid \gamma \in \mathbb{C} \right\}$$

is here contained in $\text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)})$.

Let φ be the polynomial automorphism given by $\varphi = (z_0 + z_1^2, \lambda z_1)$ with $\lambda \in \mathbb{C}^*$ and $\lambda^2 \neq 1$. A ς -lift of φ to $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ is

$$\phi = \left(z_0 + z_1^2, \lambda z_1, \lambda z_2 - \frac{z_1^3}{3} + \mu \right)$$

for some $\mu \in \mathbb{C}$. Note that

$$G = \left\{ \left(\delta z_0 + \frac{\gamma^2 - \delta}{\lambda^2 - 1} z_1 + \varepsilon, \gamma z_1 \right) \mid \delta, \gamma \in \mathbb{C}^*, \varepsilon \in \mathbb{C} \right\}$$

is contained in $\text{Cent}(\varphi, \text{Aut}(\mathbb{C}^2))$. Let us denote by G_{ς} the ς -lift of G ; a direct computation shows that

$$G_{\varsigma} = \left\{ \left(\delta z_0 + \frac{\gamma^2 - \delta}{\lambda^2 - 1} z_1 + \varepsilon, \gamma z_1, \delta \gamma z_2 - \frac{\gamma(\gamma^2 - \delta)}{3(\lambda^2 - 1)} z_1^3 - \gamma \varepsilon z_1 + \beta \right) \mid \delta, \gamma \in \mathbb{C}^*, \beta, \varepsilon \in \mathbb{C} \right\}$$

The inclusion $G_{\varsigma} \cap \text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)}) \subsetneq G_{\varsigma}$ is strict; indeed

$$G_{\varsigma} \cap \text{Cent}(\phi, \text{Aut}(\mathbb{C}^3)_{c(\omega)}) = \left\{ \left(\gamma^2 z_0 + \varepsilon, \gamma z_1, \gamma^3 z_2 - \gamma \varepsilon z_1 + \frac{\gamma^3 - 1}{\lambda - 1} \delta \right) \mid \gamma \in \mathbb{C}^*, \varepsilon \in \mathbb{C} \right\}.$$

12. NON-SIMPLICITY, TITS ALTERNATIVE

12.1. Non-simplicity. Let us recall that a **simple group** is a non-trivial group G whose only normal subgroups are $\{\text{id}\}$ and G .

Danilov proved that $\text{Aut}(\mathbb{C}^2)_\eta$ is not simple ([15]). More recently Cantat and Lamy showed that $\text{Bir}(\mathbb{P}^2)$ is not simple ([12]). As a consequence one has:

Proposition 12.1. *The groups*

$$\text{Aut}(\mathbb{C}^3)_\omega, \quad \text{Bir}(\mathbb{P}^3)_\omega, \quad \text{Aut}(\mathbb{C}^3)_{c(\omega)}, \quad [\text{Aut}(\mathbb{C}^3)_{c(\omega)}, \text{Aut}(\mathbb{C}^3)_{c(\omega)}], \quad [\text{Aut}(\mathbb{C}^3)_\omega, \text{Aut}(\mathbb{C}^3)_\omega]$$

are not simple.

Proof. Since $[\text{Aut}(\mathbb{C}^3)_{c(\omega)}, \text{Aut}(\mathbb{C}^3)_{c(\omega)}] \simeq \text{Aut}(\mathbb{C}^2)_\eta$ and $[\text{Aut}(\mathbb{C}^3)_\omega, \text{Aut}(\mathbb{C}^3)_\omega] \simeq \text{Aut}(\mathbb{C}^2)_\eta$ the first assertion follows from [15].

The exact sequence (3.1) implies in particular that there exists a morphism with a non-trivial kernel from $\text{Aut}(\mathbb{C}^3)_\omega$ into $\text{Aut}(\mathbb{C}^2)_\eta$, hence $\text{Aut}(\mathbb{C}^3)_\omega$ is not simple. A similar argument holds for $\text{Bir}(\mathbb{P}^3)_\omega$ and $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$. $\square$

The morphism

$$\text{Bir}(\mathbb{P}^3)_\omega^{\text{reg}} \longrightarrow \text{Bir}(\mathbb{P}^2)$$

that consists to take the restriction of $\phi \in \text{Bir}(\mathbb{P}^3)_\omega^{\text{reg}}$ to $\mathcal{H}_\infty$ has a non-trivial kernel; indeed

$$\phi = \left(z_0 - \left(\frac{P(z_1)}{Q(z_1)} \right)', z_1, z_2 + \frac{P(z_1)}{Q(z_1)} \right)$$

with P, Q two polynomials of degree p, q such that $p < q + 1$, is regular and induces the identity on $\mathcal{H}_\infty$. In particular one gets the following statement:

Proposition 12.2. *The group $\text{Bir}(\mathbb{P}^3)_\omega^{\text{reg}}$ is not simple.*

Let us consider the maps $\psi = \left(\gamma z_0^2 z_1, \frac{1}{\gamma z_0}, z_2 + z_0 z_1 \right)$ and $\phi = \left(z_0 + \frac{1}{z_1^3}, z_1, z_2 + \frac{1}{2z_1^2} \right)$. One can check that ψ belongs to $\text{Bir}(\mathbb{P}^3)_\omega \setminus \text{Bir}(\mathbb{P}^3)_\omega^{\text{reg}}$ whereas ϕ is in $\text{Bir}(\mathbb{P}^3)_\omega^{\text{reg}}$. A direct computation shows that $\psi^{-1}\phi\psi$ blows down $\mathcal{H}_\infty$ onto $\mathbf{e}_3$. Hence one can state:

Proposition 12.3. *The subgroup $\text{Bir}(\mathbb{P}^3)_\omega^{\text{reg}}$ of $\text{Bir}(\mathbb{P}^3)_\omega$ is not normal.*

12.2. The Tits alternative. The derived series of a group G is defined as follows

$$D_0(G) = G, \quad D_1(G) = [G, G], \quad \dots, \quad D_{n+1}(G) = [G, D_n(G)]$$

The group G is **solvable** if there exists an integer k such that $D_k(G) = \{\text{id}\}$. The least ℓ such that $D_\ell = \{\text{id}\}$ is called the **derived length** of G .

A group G satisfies the **Tits alternative** if any finitely generated subgroup of G contains either a non-abelian free group, or a solvable subgroup of finite index. This alternative has been established by Tits for linear groups $\text{GL}(n; \mathbb{k})$ for any field $\mathbb{k}$, but is not satisfied by the group of diffeomorphisms of real manifolds ([22]). Nevertheless Lamy proves that the group of polynomial automorphisms of $\text{Aut}(\mathbb{C}^2)$ satisfies the Tits alternative ([29]), so does Cantat for the group of birational maps of a complex, compact, kähler surface (see [11]). Note that the automorphisms groups of complex, compact, kähler manifolds of any dimension also satisfies Tits alternative ([11, 30]).

Theorem 12.4. *The groups $\text{Aut}(\mathbb{C}^3)_\omega$, $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ and $\text{Bir}(\mathbb{P}^3)_\omega$ satisfy the Tits alternative.*

Proof. Let G be a finitely generated subgroup of $\text{Bir}(\mathbb{P}^3)_{\omega}$. Set

$$G_0 = \zeta(G) \subset \text{Bir}(\mathbb{P}^2)_{\eta}$$

Since $\text{Bir}(\mathbb{P}^2)_{\eta}$ is a subgroup of $\text{Bir}(\mathbb{P}^2)$ that satisfies the Tits alternative, either G_0 contains a non-abelian free group, or a solvable subgroup of finite index.

Assume first that G_0 contains two elements f and h such that $\langle f, h \rangle \simeq \mathbb{Z} * \mathbb{Z}$. Let us denote by F , resp. H a lift of f , resp. h in $\text{Bir}(\mathbb{P}^3)$. Suppose that there exists a non-trivial word M such that $M(F, H) = \{\text{id}\}$. As ζ is a morphism, one gets that $M(f, h) = \{\text{id}\}$: contradiction.

Suppose now that up to finite index G_0 is solvable, and let ℓ be its derived length; in particular $D_{\ell}(G_0) = \{\text{id}\}$ and $D_{\ell}(G)$ belongs to $\ker \zeta$. Since from Proposition 4.1

$$\ker \zeta = Z(\text{Bir}(\mathbb{P}^3)_{\omega}) = \{(z_0, z_1, z_2 + \beta) \mid \beta \in \mathbb{C}\}$$

one gets $D_{\ell+1}(G) = \{\text{id}\}$. □

13. NON-CONJUGATE ISOMORPHIC GROUPS

Let us denote by v_1 the trivial embedding from $(\text{Aut}(\mathbb{C}^2)_{\eta} | 0)$ into $\text{Aut}(\mathbb{C}^3)$

$$v_1: (\text{Aut}(\mathbb{C}^2)_{\eta} | 0) \hookrightarrow \text{Aut}(\mathbb{C}^3), \quad (\phi_0, \phi_1) \mapsto (\phi_0, \phi_1, z_2)$$

and by v_2 the trivial embedding from $\text{Bir}(\mathbb{P}^2)$ into $\text{Bir}(\mathbb{P}^3)$

$$v_2: \text{Bir}(\mathbb{P}^2) \hookrightarrow \text{Bir}(\mathbb{P}^3), \quad (\phi_1, \phi_2) \mapsto (z_0, \phi_1, \phi_2).$$

Despite $\text{im } v_1$ (resp. $\text{im } v_2$) is isomorphic to $\text{im } \zeta$ (resp. $\text{im } \mathcal{K}$) one has the following statement:

Proposition 13.1. *The image of v_1 (resp. v_2) is not $\text{Aut}(\mathbb{C}^3)$ -conjugate (resp. $\text{Bir}(\mathbb{P}^3)$ -conjugate) to a subgroup of $\text{Aut}(\mathbb{C}^3)_{c(\omega)}$ (resp. $\text{Bir}(\mathbb{P}^3)_{c(\omega)}$).*

Proof. Let us assume that there exists ψ in $\text{Aut}(\mathbb{C}^3)$ (resp. $\text{Bir}(\mathbb{P}^3)$) such that for any $\phi = (\phi_0, \phi_1)$ (resp. $\phi = (\phi_1, \phi_2)$) in $\text{Aut}(\mathbb{C}^2)$ (resp. $\text{Bir}(\mathbb{P}^2)$) the map $\psi v_1(\phi) \psi^{-1}$ (resp. $\psi v_2(\phi) \psi^{-1}$) is a contact polynomial automorphism (resp. contact birational map); as a result $v_1(\phi)$ (resp. $v_2(\phi)$) preserves a polynomial form $\Theta = \text{Ad}z_0 + \text{Bdz}_1 + \text{Cdz}_2$. Looking at the restriction to any hyperplane $z_2 = \lambda$ (resp. $z_0 = \lambda$) for λ generic one gets that all the ϕ preserve the foliation given by $\Theta|_{z_2=\lambda}$ (resp. $\Theta|_{z_0=\lambda}$): contradiction. □

Part 4. Appendix: Automorphisms group of $\text{Aut}(\mathbb{C}^2)_{\eta}$

As we recalled $\text{Aut}(\mathbb{C}^2)$ is generated by J_2 and Aff_2 . More precisely $\text{Aut}(\mathbb{C}^2)$ has a structure of amalgamated product ([27])

$$\text{Aut}(\mathbb{C}^2) = J_2 *_{J_2 \cap \text{Aff}_2} \text{Aff}_2;$$

this is also the case for $\text{Aut}(\mathbb{C}^2)_{\eta}$ ([21, Proposition 9])

$$\text{Aut}(\mathbb{C}^2)_{\eta} = (J_2)_{\eta} *_{(J_2)_{\eta} \cap (\text{Aff}_2)_{\eta}} (\text{Aff}_2)_{\eta}$$

Following [16] we prove that:

Theorem 13.2. *The group $\text{Aut}(\text{Aut}(\mathbb{C}^2)_{\eta})$ is generated by the automorphisms of the field $\mathbb{C}$ and the group of $\text{Aut}(\mathbb{C}^2)$ -inner automorphisms.*

Idea of the Proof. Let us set $\mathcal{G} = \text{Aut}(\mathbb{C}^2)_{\eta}$. One can follow [16] and prove that if ϕ is an automorphism of $\mathcal{G}$, then

- $\phi((J_2)_{\eta}) = (J_2)_{\eta}$ up to conjugacy by an element of $\text{Aut}(\mathbb{C}^2)$ ([16, Proposition 4.4]);

- for any integer k if $\mathcal{R} = \cup_{n \leq k} \langle (\beta z_0, \frac{z_1}{\beta}) \mid \beta \text{ } n\text{-th root of unity} \rangle$, then there exists ψ in $(J_2)_\eta$ such that $\varphi(\mathcal{R}) = \psi \mathcal{R} \psi^{-1}$. So one can suppose that $\varphi((J_2)_\eta) = (J_2)_\eta$ and $\varphi(\mathcal{R}) = \mathcal{R}$ (see [16, Proposition 4.4]);
- set $D_\eta = \{(\beta z_0, z_1/\beta) \mid \beta \in \mathbb{C}^*\}$ one can show that conjugating ϕ by an element of $(J_2)_\eta$ one has $\varphi((J_2)_\eta) = (J_2)_\eta$ and $\varphi(D_\eta) = D_\eta$.
- set

$$T_1 = \{(z_0 + \beta, z_1) \mid \beta \in \mathbb{C}\}, \quad T_2 = \{(z_0, z_1 + \beta) \mid \beta \in \mathbb{C}\}$$

and

$$T = \{(z_0 + \gamma, z_1 + \beta) \mid \gamma, \beta \in \mathbb{C}\}$$

Since $T_1 \subset [(J_2)_\eta, (J_2)_\eta], [(J_2)_\eta, (J_2)_\eta]$, then $T_1 \subset \{(z_0 + P(z_1), z_1) \mid P \in \mathbb{C}[z_1]\}$. As

$$\forall n \in \mathbb{N}, \forall \beta \in \mathbb{C} \quad \left(\frac{z_0}{n}, nz_1\right) (z_0 + \beta, z_1)^n \left(nz_0, \frac{z_1}{n}\right) = (z_0 + \beta, z_1)$$

and $\varphi(D_\eta) = D_\eta$, one gets

$$\forall n \in \mathbb{N}, \forall \beta \in \mathbb{C} \quad \varphi\left(\frac{z_0}{n}, nz_1\right) \varphi(z_0 + \beta, z_1)^n \varphi\left(nz_0, \frac{z_1}{n}\right) = \varphi(z_0 + \beta, z_1)$$

that is

$$\forall n \in \mathbb{N} \quad \left(\frac{z_0}{\delta}, \delta z_1\right) (z_0 + nP(z_1), z_1)^n \left(\delta z_0, \frac{z_1}{\delta}\right) = (z_0 + P(z), z_1)$$

so $P(z_1) = \frac{n}{\delta} P\left(\frac{z_1}{\delta}\right)$. The polynomial P is non-zero hence $n = \delta$ and P is a constant. Therefore $\varphi(T_1) \subset T_1$.

The groups T_1 and T_2 commute, that's why

$$\varphi(T_2) \subset \{(z_0 + P(z_1), z_1 + \beta) \mid P \in \mathbb{C}[z_1], \beta \in \mathbb{C}\}$$

The relation

$$\left(\frac{z_0}{n}, nz_1\right) (z_0, z_1 + \beta) \left(nz_0, \frac{z_1}{n}\right) = (z_0, z_1 + \beta)^n$$

true for any integer n and for any β in $\mathbb{C}$ implies that $\varphi(T_2) \subset T_2$. The group T being a maximal abelian subgroup of $\mathcal{G}$, one has $\varphi(T) = T$ and $\varphi(T_i) = T_i$.

- There exist ξ_1, ξ_2 two additive morphisms and ζ a multiplicative one such that

$$\varphi(z_0 + \gamma, z_1 + \beta) = (z_0 + \xi_1(\gamma), z_1 + \xi_2(\beta)) \quad \& \quad \varphi\left(\gamma z_0, \frac{z_1}{\gamma}\right) = \left(\zeta(\gamma) z_0, \frac{z_1}{\zeta(\gamma)}\right)$$

The statement follows from [16, Proposition 1.4].

□

REFERENCES

- [1] M. Alberich-Carramiñana. *Geometry of the plane Cremona maps*, volume 1769 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2002.
- [2] A. Banyaga. Sur le groupe des difféomorphismes symplectiques. *C. R. Acad. Sci. Paris Sér. A*, 278:1343–1344, 1974.
- [3] A. Banyaga. Sur le groupe des difféomorphismes qui préservent une forme de contact régulière. *C. R. Acad. Sci. Paris Sér. A-B*, 281(16):Aii, A707–A709, 1975.
- [4] A. Banyaga. The group of diffeomorphisms preserving a regular contact form. In *Topology and algebra (Proc. Colloq., Eidgenöss. Tech. Hochsch., Zurich, 1977)*, volume 26 of *Monograph. Enseign. Math.*, pages 47–53. Univ. Genève, Geneva, 1978.
- [5] L. Bayle and A. Beauville. Birational involutions of $\mathbf{P}^2$. *Asian J. Math.*, 4(1):11–17, 2000. Kodaira's issue.

- [6] E. Bertini. Ricerche sulle trasformazioni univoche involutorie nel piano. *Annali di Mat.*, 8:244–286, 1877.
- [7] J. Blanc. Symplectic birational transformations of the plane. *Osaka J. Math.*, 50(2):573–590, 2013.
- [8] A. M. Bonifant and J. E. Fornæss. Growth of degree for iterates of rational maps in several variables. *Indiana Univ. Math. J.*, 49(2):751–778, 2000.
- [9] M. Brunella. Minimal models of foliated algebraic surfaces. *Bull. Soc. Math. France*, 127(2):289–305, 1999.
- [10] S. Cantat. *Dynamique des automorphismes des surfaces complexes compactes*. PhD thesis, École Normale Supérieure de Lyon, 1999.
- [11] S. Cantat. Morphisms between Cremona groups, and characterization of rational varieties. *Compos. Math.*, 150(7):1107–1124, 2014.
- [12] S. Cantat and S. Lamy. Normal subgroups in the Cremona group. *Acta Math.*, 210(1):31–94, 2013. With an appendix by Yves de Cornulier.
- [13] D. Cerveau and J. Déserti. *Transformations birationnelles de petit degré*, volume 19 of *Cours Spécialisés*. Société Mathématique de France, Paris, 2013.
- [14] D. Cerveau and J.-F. Mattei. *Formes intégrables holomorphes singulières*, volume 97 of *Astérisque*. Société Mathématique de France, Paris, 1982. With an English summary.
- [15] V. I. Danilov. Non-simplicity of the group of unimodular automorphisms of an affine plane. *Mat. Zametki*, 15:289–293, 1974.
- [16] J. Déserti. Sur le groupe des automorphismes polynomiaux du plan affine. *J. Algebra*, 297(2):584–599, 2006.
- [17] J. Déserti. Sur les automorphismes du groupe de Cremona. *Compos. Math.*, 142(6):1459–1478, 2006.
- [18] J. Diller and J.-L. Lin. Rational surface maps with invariant meromorphic two forms.
- [19] K. Frantzen and T. Peternell. On the bimeromorphic geometry of compact complex contact threefolds. In *Classification of algebraic varieties*, EMS Ser. Congr. Rep., pages 277–288. Eur. Math. Soc., Zürich, 2011.
- [20] S. Friedland and J. Milnor. Dynamical properties of plane polynomial automorphisms. *Ergodic Theory Dynam. Systems*, 9(1):67–99, 1989.
- [21] J.-P. Furter and S. Lamy. Normal subgroup generated by a plane polynomial automorphism. *Transform. Groups*, 15(3):577–610, 2010.
- [22] É. Ghys. Groups acting on the circle. *Enseign. Math. (2)*, 47(3–4):329–407, 2001.
- [23] M. Gizatullin. Klein’s conjecture for contact automorphisms of the three-dimensional affine space. *Michigan Math. J.*, 56(1):89–98, 2008.
- [24] M. H. Gizatullin and V. I. Danilov. Automorphisms of affine surfaces. I. *Izv. Akad. Nauk SSSR Ser. Mat.*, 39(3):523–565, 703, 1975.
- [25] E. L. Ince. *Ordinary Differential Equations*. Dover Publications, New York, 1944.
- [26] J. P. Jouanolou. *Équations de Pfaff algébriques*, volume 708 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979.
- [27] H. W. E. Jung. Über ganze birationale Transformationen der Ebene. *J. Reine Angew. Math.*, 184:161–174, 1942.
- [28] H. Kraft and I. Stampfli. On automorphisms of the affine cremona group. *Ann. Inst. Fourier (Grenoble)*, 63(3):1535–1543, 2013.
- [29] S. Lamy. L’alternative de Tits pour $\text{Aut}[\mathbb{C}^2]$. *J. Algebra*, 239(2):413–437, 2001.
- [30] K. Oguiso. Tits alternative in hyperkähler manifolds. *Math. Res. Lett.*, 13:307–316, 2006.
- [31] A. van den Essen. *Polynomial automorphisms and the Jacobian conjecture*, volume 190 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2000.

IRMAR, UMR 6625 DU CNRS, UNIVERSITÉ DE RENNES 1, 35042 RENNES, FRANCE.

E-mail address: dominique.cerveau@univ-rennes1.fr

UNIVERSITÉ PARIS DIDEROT, SORBONNE PARIS CITÉ, INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, UMR 7586, CNRS, SORBONNE UNIVERSITÉS, UPMC UNIV PARIS 06, F-75013 PARIS, FRANCE.

E-mail address: deserti@math.univ-paris-diderot.fr